

ВН



WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

(51) International Patent Classification ⁷ : C12N 9/00, C07K 14/00	A3	(11) International Publication Number: WO 00/21986
		(43) International Publication Date: 20 April 2000 (20.04.00)
(21) International Application Number: PCT/US99/23315 (22) International Filing Date: 6 October 1999 (06.10.99) (30) Priority Data: 09/169,289 9 October 1998 (09.10.98) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 09/169,289 (CIP) Filed on 9 October 1998 (09.10.98)		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> (88) Date of publication of the international search report: 13 July 2000 (13.07.00)
(71) Applicant (<i>for all designated States except US</i>): INCYTE PHARMACEUTICALS, INC. [US/US]; 3174 Porter Drive, Palo Alto, CA 94304 (US). (72) Inventors; and (75) Inventors/Applicants (<i>for US only</i>): WALKER, Michael, G. [CA/US]; Unit 80, 1050 Borregas Avenue, Sunnyvale, CA 94089 (US). VOLKMUTH, Wayne [US/US]; 783 Roble Avenue, #1, Menlo Park, CA 94025 (US). KLINGLER, Tod, M. [US/US]; 28 Dover Court, San Carlos, CA 94070 (US). (74) Agents: BILLINGS, Lucy, J. et al.; Incyte Pharmaceuticals, Inc., 3174 Porter Drive, Palo Alto, CA 94304 (US).		
(54) Title: MATRIX-REMODELING GENES		
(57) Abstract <p>The invention provides novel matrix-remodeling genes and polypeptides encoded by those genes. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating or preventing diseases associated with matrix remodeling.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/23315

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N9/00 C07K14/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
------------	--	-----------------------

A	MATRISIAN, L. M.: "METALLOPROTEINASES AND THEIR INHIBITORS IN MATRIX REMODELING" TRENDS IN GENETICS, vol. 6, no. 4, 1990, pages 121-125, XP000644304 * figure 2; page 123 *	1-12
---	--	------

A	YE, S. ET AL.: "MATRIX METALLOPROTEINASES: ..." CLINICAL SCIENCE, vol. 94, February 1998 (1998-02), pages 103-110, XP000857426 * whole disclosure *	1-12
---	--	------

-/--

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

Z document member of the same patent family

Date of the actual completion of the international search

27 January 2000

Date of mailing of the international search report

03.05.00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax (+31-70) 340-3016

Authorized officer

HERMANN R.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/23315

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE EMBL/GENBANK [Online] ID HSZZ74011; AC AA368885, 18 April 1997 (1997-04-18) ADAMS, M.D. ET AL.: "EST80295 Placenta I homo sapiens cDNA" XP002128855 abstract</p> <p>---</p>	2-8
X	<p>DATABASE EMBL/GENBANK [Online] ID HSW4574; AC W92457, 18 July 1996 (1996-07-18) HILLIER, L. ET AL.: "EST; Soares fetal heart NbHH19W homo sapiens cDNA clone" XP002128856 abstract</p> <p>-----</p>	2-8

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/23315

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 10-12
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 10-12 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.: 2,7
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
See FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

See additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-3,5-7,9,10,12 (all partially)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 2,7

Claims 2 and 7 are -inter alia- drafted to compounds comprising as few as 18 nucleotides / 6 amino acids, which are neither defined by their exact structure, not by their exact location in the parent molecule.

Moreover, claim 2 covers

- polynucleotides of completely undefined length that hybridize to said short sequence (or the parent), and
- sequences which are complementary to the almost undefined sequences mentioned above.

Said vague structural definitions are not sufficient for a reasonable search.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: 1-3,5-7,9,10,12 (all partially)

A polynucleotide having the SEQ ID NO 1; subject-matter of claims 2,3,5,6,9,12 relating to said polynucleotide

2. Claims: 1-12, all partially

Polypeptide with the SEQ ID NO. 21, encoding polynucleotides, and related subject-matter

3. Claims: 1-12, all partially

Polypeptide with the SEQ ID NO. 22, encoding polynucleotides, and related subject-matter

4. Claims: 1-12, all partially

Polypeptide with the SEQ ID NO. 23, encoding polynucleotides, and related subject-matter

5. Claims: 1-3,5-7,9,10,12 (all partially)

INVENTIONS 5-20:
Polynucleotides having the SEQ ID NOs. 3-5,7-10,12-20; subject-matter of claims 2,3,5,6,9,12 relating to said polynucleotide.

6. Claims: 1,3 (partially)

INVENTION 21:
Further polynucleotides and encoded polypeptides of claim 1 (PLEASE NOTE: This subject has been added for completeness only. It is considered as being at least partially anticipated (infra); Further non-unity objections may arise upon payment for subject 21)

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07K 14/00	A2	(11) International Publication Number: WO 00/21986 (43) International Publication Date: 20 April 2000 (20.04.00)
(21) International Application Number: PCT/US99/23315 (22) International Filing Date: 6 October 1999 (06.10.99) (30) Priority Data: 09/169,289 9 October 1998 (09.10.98) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 09/169,289 (CIP) Filed on 9 October 1998 (09.10.98) (71) Applicant (for all designated States except US): INCYTE PHARMACEUTICALS, INC. [US/US]; 3174 Porter Drive, Palo Alto, CA 94304 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): WALKER, Michael, G. [CA/US]; Unit 80, 1050 Borregas Avenue, Sunnyvale, CA 94089 (US). VOLKMUTH, Wayne [US/US]; 783 Roble Avenue, #1, Menlo Park, CA 94025 (US). KLINGLER, Tod, M. [US/US]; 28 Dover Court, San Carlos, CA 94070 (US). (74) Agents: BILLINGS, Lucy, J. et al.; Incyte Pharmaceuticals, Inc., 3174 Porter Drive, Palo Alto, CA 94304 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: MATRIX-REMODELING GENES (57) Abstract The invention provides novel matrix-remodeling genes and polypeptides encoded by those genes. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating or preventing diseases associated with matrix remodeling.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

MATRIX-REMODELING GENES

TECHNICAL FIELD

5 The invention relates to novel matrix-remodeling genes identified by their
coexpression with known matrix-remodeling genes. The invention also relates to the use of
these biomolecules in diagnosis, prognosis, prevention, treatment, and evaluation of therapies
for diseases, particularly diseases associated with matrix-remodeling such as cancer,
cardiomyopathy, arthritis, angiogenesis, diabetic necrosis, atherosclerosis, fibrosis, and
10 ulceration.

BACKGROUND OF THE INVENTION

Matrix remodeling is associated with the construction, destruction, and reorganization
of extracellular matrix components and is essential in normal cellular functions and also in
many disease processes. These disease processes include metastatic cancer, cardiomyopathy,
15 arthritis, angiogenesis, diabetic necrosis, atherosclerosis, fibrosis, and ulceration (Alexander
and Werb (1991) In: Cell Biology of Extracellular Matrix, Plenum Press, New York NY, pp.
255-302; Schuppan et al. (1993) In: Extracellular Matrix, Marcel Dekker, New York NY, pp.
201-254; Zvibel and Kraft (1993) In: Extracellular Matrix, Marcel Dekker, New York NY, pp.
559-580; Shanahan et al. (1994) J Clin Invest 93:2393-402; Kielty and Shuttleworth (1995) Int
20 J Biochem Cell Biol 27:747-60; Bitar and Labbad (1996) J Surg Res 61:113-9; Dourado et al.
(1996) Osteoarthritis Cartilage 4:187-96; Grant et al. (1996) Regul. Pept. 67:137-44; Gunja-
Smith et al. (1996) Am J Pathol 148:1639-48; Alcolado et al. (1997) Clin. Sci 92:103-12; Cs-
Szabo et al. (1997) Arthritis Rheum 40:1037-45; Hayward and Brock (1997) Hum Mutat
10:415-23; Ledda et al. (1997) J Invest Dermatol 108:210-4; Hayashido et al. (1998) Int J
25 Cancer 75:654-8; Ito et al. (1998) Kidney Int 53:853-61; Nelson et al. (1998) Cancer Res
58:232-6).

Many genes that participate in and regulate matrix remodeling are known, but many
remain to be identified. Identification of currently unknown genes will provide new
diagnostic and therapeutic targets. In addition, these genes will provide new opportunities for
30 therapeutic tissue engineering--the use of drugs or biologicals to direct the creation of new
tissues such as skin, pancreas, or liver that can replace tissues lost to disease or trauma.

The present invention provides new compositions that are useful for diagnosis, prognosis, treatment, prevention, and evaluation of therapies for diseases associated with matrix remodeling. We have implemented a method for analyzing gene expression patterns and have identified 20 novel matrix-remodeling genes by their coexpression with known
5 matrix-remodeling genes.

SUMMARY OF THE INVENTION

In one aspect, the invention provides for a substantially purified polynucleotide comprising a gene that is coexpressed with one or more known matrix-remodeling genes in a plurality of biological samples. Preferably, each known matrix-remodeling gene is selected
10 from the group consisting of osteonectin (BM-40), chondroitin/dermatan sulfate proteoglycans (C/DSPG), collagen I, II, III, and IV, connective tissue growth factor (CTGF), fibrillin, fibronectins, fibronectin receptor (fibr-r), fibulin 1, heparan sulfate proteoglycans (HSPG), extracellular matrix protein (hevin), insulin-like growth factor 1 (IGF 1), insulin-like growth factor binding protein (IGFBP), laminin, lumican, matrix Gla protein (MGP), matrix
15 metalloproteases (MMP), and tissue inhibitors of matrix metalloproteinase 1, 2, and 3 (TIMP 1, 2, and 3). Preferred embodiments are (a) a polynucleotide sequence selected from the group consisting of SEQ ID NOs:1-20; (b) a polynucleotide sequence which encodes a polypeptide sequence of SEQ ID NOs:21, 22, and 23; (c) a polynucleotide sequence having at least 70% identity to the polynucleotide sequence of (a) or (b); (d) a polynucleotide
20 sequence comprising at least 18 sequential nucleotides of the polynucleotide sequence of (a), (b), or (c); (e) a polynucleotide which hybridizes under stringent conditions to the polynucleotide of (a), (b), (c), or (d); or (f) a polynucleotide sequence which is complementary to the polynucleotide sequence of (a), (b), (c), (d) or (e). Furthermore, the invention provides an expression vector comprising any of the above described
25 polynucleotides and host cells comprising the expression vector. Still further, the invention provides a method for treating or preventing a disease or condition associated with the altered expression of a gene that is coexpressed with one or more known matrix-remodeling genes in a sample comprising administering to a subject in need the above-described polynucleotides in an amount effective for treating or preventing said disease.

30 In a second aspect, the invention provides a substantially purified polypeptide comprising the gene product of a gene that is coexpressed with one or more known matrix-

remodeling genes in a plurality of biological samples. The known matrix-remodeling gene may be selected from the group consisting of osteonectin (BM-40), chondroitin/dermatan sulfate proteoglycans (C/DSPG), collagen I, II, II, and IV, connective tissue growth factor (CTGF), fibrillin, fibronectins, fibronectin receptors (fibr-r), fibulin 1, heparan sulfate
5 proteoglycans (HSPG), extracellular matrix protein (hevin), insulin-like growth factor 1 (IGF 1), insulin-like growth factor binding protein (IGFBP), laminin, lumican, matrix Gla protein (MGP), matrix metalloproteases (MMP), and tissue inhibitors of matrix metalloproteinase 1, 2, and 3 (TIMP 1, 2, and 3). Preferred embodiments are polypeptides comprising (a) the polypeptide sequence of SEQ ID NO:21, 22, or 23; (b) a polypeptide sequence having at least
10 85% identity to the polypeptide sequence of (a); and (c) a polypeptide sequence comprising at least 6 sequential amino acids of the polypeptide sequence of (a) or (b). Additionally, the invention provides antibodies that bind specifically to any of the above described polypeptides and a method for treating or preventing a disease or condition associated with the altered expression of a gene that is coexpressed with one or more known matrix-remodeling genes in
15 a sample comprising administering to a subject in need such an antibody in an amount effective for treating or preventing said disease.

In another aspect, the invention provides a pharmaceutical composition comprising the polynucleotide of claim 2 or the polypeptide of claim 3 in conjunction with a suitable pharmaceutical carrier or a method for treating or preventing a disease or condition associated
20 with the altered expression of a gene that is coexpressed with one or more known matrix-remodeling genes in a sample comprising administering to a subject in need such compositioning in an amount effective for treating or preventing said disease.

In yet a further aspect, the invention provides a method for diagnosing a disease or condition associated with the altered expression of a gene that is coexpressed with one or more
25 known matrix-remodeling genes in a sample, wherein each known matrix-remodeling gene is selected from the group consisting of osteonectin (BM-40), chondroitin/dermatan sulfate proteoglycans (C/DSPG), collagen I, II, II, and IV, connective tissue growth factor (CTGF), fibrillin, fibronectins, fibronectin receptor (fibr-r), fibulin 1, heparan sulfate proteoglycans (HSPG), extracellular matrix protein (hevin), insulin-like growth factor 1 (IGF 1), insulin-like
30 growth factor binding protein (IGFBP), laminin, lumican, matrix Gla protein (MGP), matrix metalloproteases (MMP), and tissue inhibitors of matrix metalloproteinase 1, 2, and 3 (TIMP

1, 2, and 3). The method comprises the steps of (a) providing the sample comprising one of more of said coexpressed genes; (b) hybridizing the polynucleotide of the coexpressed genes under conditions effective to form one or more hybridization complexes; and (c) detecting the hybridization complexes, wherein the altered level of one or more of the hybridization
5 complexes in a diseased sample compared with the level of hybridization complexes in a non-diseased sample correlates with the presence of the disease or condition in the sample.

BRIEF DESCRIPTION OF THE SEQUENCE LISTING

The Sequence Listing provides exemplary matrix-remodeling-associated sequences including polynucleotide sequences, SEQ ID NOs:1-20, and polypeptide sequences, SEQ ID
10 NOs:21-23. Each sequence is identified by a sequence identification number (SEQ ID NO) and by the Incyte Clone number from which the sequence was first identified.

DESCRIPTION OF THE INVENTION

It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include the plural reference unless the context clearly dictates otherwise.
15 Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

DEFINITIONS

"NSEQ" refers generally to a polynucleotide sequence of the present invention,
20 including SEQ ID NOs:1-20. "PSEQ" refers generally to a polypeptide sequence of the present invention, including SEQ ID NOs:21-23.

A "variant" refers to either a polynucleotide or a polypeptide whose sequence diverges from SEQ ID NOs:1-20 or SEQ ID NOs:21-23, respectively. Polynucleotide sequence divergence may result from mutational changes such as deletions, additions, and substitutions
25 of one or more nucleotides; it may also occur because of differences in codon usage. Each of these types of changes may occur alone, or in combination, one or more times in a given sequence. Polypeptide variants include sequences that possess at least one structural or functional characteristic of SEQ ID NOs:21-23.

A "fragment" can refer to a nucleic acid sequence that is preferably at least 20 nucleic
30 acids in length, more preferably 40 nucleic acids, and most preferably 60 nucleic acids in length, and encompasses, for example, fragments consisting of nucleic acids 1-50 or 200-500

of SEQ ID NOs:1-20. A "fragment" can also refer to polypeptide sequences which are preferably at least 5 to about 15 amino acids in length, most preferably at least 10 amino acids long, and which retain some biological activity or immunological activity of, for example, a sequence selected from SEQ ID NOs:21-23.

- 5 "Gene" or "gene sequence" refers to the partial or complete coding sequence of a transcript. The term also refers to sequences corresponding to 5' or 3' untranslated regions or 5' or 3' untranslated regions including partial or complete coding sequences of a gene. Typically, the novel gene sequences may or may not be homologous to annotated sequences found in public or private databases. The gene may be in a sense or antisense
10 (complementary) orientation.

"Known matrix-remodeling gene" refers to a gene sequence which has been previously identified as useful in the diagnosis, treatment, prognosis, or prevention of diseases associated with matrix remodeling. Typically, this means that the known matrix-remodeling gene is expressed at higher levels in tissue abundant in known matrix-remodeling transcripts when
15 compared with other tissue.

"Matrix-remodeling gene" refers to a gene sequence whose expression pattern is similar to that of the known matrix-remodeling genes and which are useful in the diagnosis, treatment, prognosis, or prevention of diseases associated with matrix remodeling. The gene sequences can also be used in the evaluation of therapies for cancer.

- 20 "Substantially purified" refers to a nucleic acid or an amino acid sequence that is removed from its natural environment and is isolated or separated, and is at least about 60% free, preferably about 75% free, and most preferably about 90% free from other components with which it is naturally present.

THE INVENTION

- 25 The present invention encompasses a method for identifying biomolecules that are associated with a specific disease, regulatory pathway, subcellular compartment, cell type, tissue type, or species. In particular, the method identifies gene sequences useful in diagnosis, prognosis, treatment, prevention, and evaluation of therapies for diseases associated with matrix-remodeling, particularly, cancer, cardiomyopathy, arthritis, angiogenesis, diabetic
30 necrosis, atherosclerosis, fibrosis, and ulceration.

The method provides first identifying polynucleotides that are expressed in a plurality

of cDNA libraries. The identified polynucleotides include genes of known function, genes known to be specifically expressed in a specific disease process, subcellular compartment, cell type, tissue type, or species. Additionally, the polynucleotides include genes of unknown function. The expression patterns of the known genes are then compared with those of the
5 genes of unknown function to determine whether a specified coexpression probability threshold is met. Through this comparison, a subset of the polynucleotides for unknown function genes having a high coexpression probability with the known genes can be identified. The high coexpression probability correlates with a particular coexpression probability threshold which is less than 0.001, and more preferably less than 0.00001.

10 The polynucleotides originate from cDNA libraries derived from a variety of sources including, but not limited to, eukaryotes such as human, mouse, rat, dog, monkey, plant, and yeast and prokaryotes such as bacteria and viruses. These polynucleotides can also be selected from a variety of sequence types including, but not limited to, expressed sequence tags (ESTs), assembled polynucleotide sequences, full length gene coding regions, introns,
15 regulatory sequences, 5' untranslated regions, and 3' untranslated regions. To have statistically significant analytical results, the polynucleotides need to be expressed in at least three cDNA libraries.

The cDNA libraries used in the coexpression analysis of the present invention can be obtained from blood vessels, heart, blood cells, cultured cells, connective tissue, epithelium,
20 islets of Langerhans, neurons, phagocytes, biliary tract, esophagus, gastrointestinal system, liver, pancreas, fetus, placenta, chromaffin system, endocrine glands, ovary, uterus, penis, prostate, seminal vesicles, testis, bone marrow, immune system, cartilage, muscles, skeleton, central nervous system, ganglia, neuroglia, neurosecretory system, peripheral nervous system, bronchus, larynx, lung, nose, pleurus, ear, eye, mouth, pharynx, exocrine glands, bladder,
25 kidney, ureter, and the like. The number of cDNA libraries selected can range from as few as 20 to greater than 10,000. Preferably, the number of the cDNA libraries is greater than 500.

In a preferred embodiment, gene sequences are assembled to reflect related sequences, such as assembled sequence fragments derived from a single transcript. Assembly of the polynucleotide sequences can be performed using sequences of various types including, but
30 not limited to, ESTs, extensions, or shotgun sequences. In a most preferred embodiment, the polynucleotide sequences are derived from human sequences that have been assembled using

the algorithm disclosed in "Database and System for Storing, Comparing and Displaying Related Biomolecular Sequence Information", Lincoln et al., Serial No:60/079,469, filed March 26, 1998, incorporated herein by reference.

Experimentally, differential expression of the polynucleotides can be evaluated by
5 methods including, but not limited to, differential display by spatial immobilization or by gel electrophoresis, genome mismatch scanning, representational difference analysis, and transcript imaging. Additionally, differential expression can be assessed by microarray technology. These methods may be used alone or in combination.

Known matrix-remodeling genes can be selected based on the use of the genes as
10 diagnostic or prognostic markers or as therapeutic targets for diseases associated with matrix remodeling, such as cancer, cardiomyopathy, arthritis, angiogenesis, diabetic necrosis, atherosclerosis, fibrosis, and ulceration. Preferably, the known matrix-remodeling genes include osteonectin (BM-40), chondroitin/dermatan sulfate proteoglycans (C/DSPG), collagen I, II, III, and IV, connective tissue growth factor (CTGF), fibrillin, fibronectins,
15 fibronectin receptor (fibr-r), fibulin 1, heparan sulfate proteoglycans (HSPG), extracellular matrix protein (hevin), insulin-like growth factor I (IGF I), insulin-like growth factor binding protein (IGFBP), laminin, lumican, matrix Gla protein (MGP), matrix metalloproteinases (MMP), tissue inhibitors of matrix metalloproteinase 1, 2, and 3 (TIMP 1, 2, and 3), and the like.

20 The procedure for identifying novel genes that exhibit a statistically significant coexpression pattern with known matrix-remodeling genes is as follows. First, the presence or absence of a gene sequence in a cDNA library is defined: a gene is present in a cDNA library when at least one cDNA fragment corresponding to that gene is detected in a cDNA sample taken from the library, and a gene is absent from a library when no corresponding cDNA
25 fragment is detected in the sample.

Second, the significance of gene coexpression is evaluated using a probability method to measure a due-to-chance probability of the coexpression. The probability method can be the Fisher exact test, the chi-squared test, or the kappa test. These tests and examples of their applications are well known in the art and can be found in standard statistics texts (Agresti, A
30 (1990) Categorical Data Analysis, John Wiley & Sons, New York NY; Rice, JA (1988) Mathematical Statistics and Data Analysis, Duxbury Press, Pacific Grove CA). A Bonferroni

correction (Rice, supra, page 384) can also be applied in combination with one of the probability methods for correcting statistical results of one gene versus multiple other genes. In a preferred embodiment, the due-to-chance probability is measured by a Fisher exact test, and the threshold of the due-to-chance probability is set to less than 0.001, more preferably less than 0.00001.

To determine whether two genes, A and B, have similar coexpression patterns, occurrence data vectors can be generated as illustrated in Table 1, wherein a gene's presence is indicated by a one and its absence by a zero. A zero indicates that the gene did not occur in the library, and a one indicates that it occurred at least once.

Table 1. Occurrence data for genes A and B

	Library 1	Library 2	Library 3	...	Library N
gene A	1	1	0	...	0
gene B	1	0	1	...	0

For a given pair of genes, the occurrence data in Table 1 can be summarized in a 2 x 2 contingency table.

Table 2. Contingency table for co-occurrences of genes A and B

	Gene A present	Gene A absent	Total
Gene B present	8	2	10
Gene B absent	2	18	20
Total	10	20	30

Table 2 presents co-occurrence data for gene A and gene B in a total of 30 libraries. Both gene A and gene B occur 10 times in the libraries. Table 2 summarizes and presents 1) the number of times gene A and B are both present in a library, 2) the number of times gene A and B are both absent in a library, 3) the number of times gene A is present while gene B is absent, and 4) the number of times gene B is present while gene A is absent. The upper left entry is the number of times the two genes co-occur in a library, and the middle right entry is the number of times neither gene occurs in a library. The off diagonal entries are the number of times one gene occurs while the other does not. Both A and B are present eight times and absent 18 times, gene A is present while gene B is absent two times, and gene B is present while gene A

is absent two times. The probability ("p-value") that the above association occurs due to chance as calculated using a Fisher exact test is 0.0003. Associations are generally considered significant if a p-value is less than 0.01 (Agresti, supra; Rice, supra).

This method of estimating the probability for coexpression of two genes makes several
5 assumptions. The method assumes that the libraries are independent and are identically
sampled. However, in practical situations, the selected cDNA libraries are not entirely
independent because more than one library may be obtained from a single patient or tissue,
and they are not entirely identically sampled because different numbers of cDNAs may be
sequenced from each library (typically ranging from 5,000 to 10,000 cDNAs per library). In
10 addition, because a Fisher exact coexpression probability is calculated for each gene versus
41,419 other genes, a Bonferroni correction for multiple statistical tests is necessary.

Using the method of the present invention, we have identified 20 novel genes that
exhibit strong association, or coexpression, with known genes that are matrix-remodeling-
specific. These known matrix-remodeling genes include osteonectin (BM-40),
15 chondroitin/dermatan sulfate proteoglycans (C/DSPG), collagen I, II, III, and IV, connective
tissue growth factor (CTGF), fibrillin, fibronectins, fibronectin receptor (fibr-r), fibulin 1,
heparan sulfate proteoglycans (HSPG), extracellular matrix protein (hevin), insulin-like
growth factor 1 (IGF 1), insulin-like growth factor binding protein (IGFBP), laminin, lumican,
matrix Gla protein (MGP), matrix metalloproteases (MMP), and tissue inhibitors of matrix
20 metalloproteinase 1, 2, and 3 (TIMP 1, 2, and 3). The results presented in Tables 5 and 6
show that the expression of the 20 novel genes have direct or indirect association with the
expression of known matrix-remodeling genes. Therefore, the novel genes can potentially be
used in diagnosis, treatment, prognosis, or prevention of diseases associated with matrix
remodeling, or in the evaluation of therapies for diseases associated with matrix remodeling.
25 Further, the gene products of the 20 novel genes are potential therapeutic proteins and targets
of therapeutics against diseases associated with matrix remodeling.

Therefore, in one embodiment, the present invention encompasses a polynucleotide
sequence comprising the sequence of SEQ ID NOs:1-20. These 20 polynucleotides are shown
by the method of the present invention to have strong coexpression association with known
30 matrix-remodeling genes and with each other. The invention also encompasses a variant of
the polynucleotide sequence, its complement, or 18 consecutive nucleotides of a sequence

provided in the above described sequences. Variant polynucleotide sequences typically have at least about 70%, more preferably at least about 85%, and most preferably at least about 95% polynucleotide sequence identity to NSEQ.

One preferred method for identifying variants entails using NSEQ and/or PSEQ
5 sequences to search against the GenBank primate (pri), rodent (rod), and mammalian (mam), vertebrate (vrtp), and eukaryote (eukp) databases, SwissProt, BLOCKS (Bairoch et al. (1997) Nucleic Acids Res 25:217-221), PFAM, and other databases that contain previously identified and annotated motifs, sequences, and gene functions. Methods that search for primary sequence patterns with secondary structure gap penalties (Smith et al. (1992) Protein
10 Engineering 5:35-51) as well as algorithms such as BLAST (Basic Local Alignment Search Tool; Altschul (1993) J Mol Evol 36:290-300; and Altschul et al. (1990) J Mol Biol 215:403-410), BLOCKS (Henikoff and Henikoff (1991) Nucleic Acids Res 19:6565-6572), Hidden Markov Models (HMM; Eddy (1996) Cur Opin Str Biol 6:361-365; Sonnhammer et al. (1997) Proteins 28:405-420), and the like, can be used to manipulate and analyze nucleotide and
15 amino acid sequences. These databases, algorithms and other methods are well known in the art and are described in Ausubel et al. (1997; Short Protocols in Molecular Biology, John Wiley & Sons, New York NY) and in Meyers (1995; Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853).

Also encompassed by the invention are polynucleotide sequences that are capable of
20 hybridizing to SEQ ID NOs:1-20, and fragments thereof under stringent conditions. Stringent conditions can be defined by salt concentration, temperature, and other chemicals and conditions well known in the art. In particular, stringency can be increased by reducing the concentration of salt, or raising the hybridization temperature. Varying additional parameters, such as hybridization time, the concentration of detergent or solvent, and the inclusion or
25 exclusion of carrier DNA, are well known to those skilled in the art. Additional variations on these conditions will be readily apparent to those skilled in the art (Wahl and Berger (1987) Methods Enzymol 152:399-407; Kimmel (1987) Methods Enzymol 152:507-511; Ausubel supra; and Sambrook et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY).

30 NSEQ or the polynucleotide sequences encoding PSEQ can be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to

detect upstream sequences, such as promoters and regulatory elements. (See, e.g., Dieffenbach and Dveksler (1995) PCR Primer, a Laboratory Manual, Cold Spring Harbor Press, Plainview NY; Sarkar (1993) PCR Methods Applic 2:318-322; Triglia et al. (1988) Nucleic Acids Res 16:8186; Lagerstrom et al. (1991) PCR Methods Applic 1:111-119; and
5 Parker et al. (1991) Nucleic Acids Res 19:3055-306). Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto, CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences,
10 Plymouth MN) or another appropriate program, to be about 18 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

In another aspect of the invention, NSEQ or the polynucleotide sequences encoding PSEQ can be cloned in recombinant DNA molecules that direct expression of PSEQ or the
15 polypeptides encoded by NSEQ, or structural or functional fragments thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and used to express the polypeptides of PSEQ or the polypeptides encoded by NSEQ. The nucleotide sequences of the present invention can be engineered using methods
20 generally known in the art in order to alter the nucleotide sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce
25 mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

In order to express a biologically active polypeptide encoded by NSEQ, NSEQ or the polynucleotide sequences encoding PSEQ, or derivatives thereof, may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for
30 transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible

promoters, and 5' and 3' untranslated regions in the vector and in NSEQ or polynucleotide sequences encoding PSEQ. Methods which are well known to those skilled in the art may be used to construct expression vectors containing NSEQ or polynucleotide sequences encoding PSEQ and appropriate transcriptional and translational control elements. These methods
5 include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook (supra) and Ausubel (supra).

A variety of expression vector/host cell systems may be utilized to contain and express NSEQ or polynucleotide sequences encoding PSEQ. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or
10 cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (baculovirus); plant cell systems transformed with viral expression vectors, cauliflower mosaic virus (CaMV) or tobacco mosaic virus (TMV), or with bacterial expression vectors (Ti or pBR322 plasmids); or animal cell systems. The invention is not limited by the host cell employed. For long term production of
15 recombinant proteins in mammalian systems, stable expression of a polypeptide encoded by NSEQ in cell lines is preferred. For example, NSEQ or sequences encoding PSEQ can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector.

20 In general, host cells that contain NSEQ and that express PSEQ may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences. Immunological
25 methods for detecting and measuring the expression of PSEQ using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS).

Host cells transformed with NSEQ or polynucleotide sequences encoding PSEQ may
30 be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly

depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides of NSEQ or polynucleotides encoding PSEQ may be designed to contain signal sequences which direct secretion of PSEQ or polypeptides encoded by NSEQ through a prokaryotic or eukaryotic cell membrane.

5 In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" form of the protein may also be used to specify protein targeting, folding,
10 and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38), are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant NSEQ or
15 nucleic acid sequences encoding PSEQ are ligated to a heterologous sequence resulting in translation of a fusion protein containing heterologous protein moieties in any of the aforementioned host systems. Such heterologous protein moieties facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin
20 (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, hemagglutinin (HA) and monoclonal antibody epitopes..

In another embodiment, NSEQ or sequences encoding PSEQ are synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers et al. (1980) Nucleic Acids Symp Ser (7) 215-223; Horn et al. (1980) Nucleic Acids Symp Ser (7) 225-232;
25 and Ausubel, supra). Alternatively, PSEQ or a polypeptide sequence encoded by NSEQ itself, or a fragment thereof, may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solid-phase techniques (Roberge et al. (1995) Science 269:202-204). Automated synthesis may be achieved using the ABI 431A Peptide synthesizer (PE Biosystems, Foster City CA). Additionally, PSEQ or the amino acid
30 sequence encoded by NSEQ, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a polypeptide

variant.

In another embodiment, the invention provides a substantially purified polypeptide comprising the amino acid sequence selected from the group consisting of SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23 or fragments thereof.

5 DIAGNOSTICS and THERAPEUTICS

The sequences of these genes can be used in diagnosis, prognosis, treatment, prevention, and evaluation of therapies for diseases associated with matrix-remodeling, particularly cancer, cardiomyopathy, arthritis, angiogenesis, diabetic necrosis, atherosclerosis, fibrosis, and ulceration. Further, the amino acid sequences encoded by the novel genes are
10 potential therapeutic proteins and targets of anti-cancer therapeutics or for the treatment of other diseases associated with matrix remodeling.

In one preferred embodiment, the polynucleotide sequences of NSEQ or the polynucleotides encoding PSEQ are used for diagnostic purposes to investigate the altered expression of PSEQ, and to monitor regulation of the levels of mRNA or the polypeptides
15 encoded by NSEQ during therapeutic intervention. The polynucleotides may be at least 18 nucleotides long, and may be complementary RNA or DNA molecules, branched nucleic acids, or peptide nucleic acids (PNAs). Alternatively, the polynucleotides are used to detect and quantitate gene expression in samples in which expression of PSEQ or the polypeptides encoded by NSEQ are correlated with disease. Additionally, NSEQ or the polynucleotides
20 encoding PSEQ can be used to detect genetic polymorphisms associated with a disease. These polymorphisms may be detected at the transcript cDNA or genomic level.

The specificity of the probe, whether it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification (maximal, high, intermediate, or low), will determine
25 whether the probe identifies only naturally occurring sequences encoding PSEQ, allelic variants, or related sequences.

Probes may also be used for the detection of related sequences, and should preferably have at least 70% sequence identity to any of the NSEQ or PSEQ-encoding sequences.

Means for producing specific hybridization probes for DNAs encoding PSEQ include
30 the cloning of NSEQ or polynucleotide sequences encoding PSEQ into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available,

and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as ^{32}P or ^{35}S , or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, by fluorescent labels and the like. The polynucleotide sequences encoding PSEQ may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; and in microarrays utilizing fluids or tissues from patients to detect altered NSEQ expression. Such qualitative or quantitative methods are well known in the art.

10 NSEQ or the nucleotide sequences encoding PSEQ can be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantitated and compared with a standard value, typically, derived from a non-diseased sample. If the amount of signal in the patient sample is significantly
15 altered in comparison to the standard value then the presence of altered levels of nucleotide sequences of NSEQ and those encoding PSEQ in the sample indicates the presence of the associated disease. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

20 Once the presence of a disease is established and a treatment protocol is initiated, hybridization or amplification assays can be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in a healthy subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

25 The polynucleotides may be used for the diagnosis of a variety of diseases associated with matrix-remodeling including cancer such as adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary
30 glands, skin, spleen, testis, thymus, thyroid, and uterus, cardiomyopathy, arthritis, angiogenesis, diabetic necrosis, atherosclerosis, fibrosis, and ulceration.

Alternatively, the polynucleotides may be used as targets in a microarray. The microarray can be used to monitor the expression level of large numbers of genes simultaneously and to identify splice variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disease, to diagnose a disease, and to develop and monitor the activities of therapeutic agents.

In yet another alternative, polynucleotides may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. Fluorescent in situ hybridization (FISH) may be correlated with other physical chromosome mapping techniques and genetic map data. (See, e.g., Heinz-Ulrich et al. (1995) in Meyers, supra, pp. 965-968.)

10 In another embodiment, antibodies which specifically bind PSEQ may be used for the diagnosis of diseases characterized by the over-or-underexpression of PSEQ or polypeptides encoded by NSEQ. A variety of protocols for measuring PSEQ or the polypeptides encoded by NSEQ, including ELISAs, RIAs, and FACS, are well known in the art and provide a basis for diagnosing altered or abnormal levels of the expression of PSEQ or the polypeptides
15 encoded by NSEQ. Standard values for PSEQ expression are established by combining body fluids or cell extracts taken from healthy subjects, preferably human, with antibody to PSEQ or a polypeptide encoded by NSEQ under conditions suitable for complex formation. The amount of complex formation may be quantitated by various methods, preferably by photometric means. Quantities of PSEQ or the polypeptides encoded by NSEQ expressed in
20 disease samples from, for example, biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing or monitoring disease. Alternatively, one may use competitive drug screening assays in which neutralizing antibodies capable of binding PSEQ or the polypeptides encoded by NSEQ specifically compete with a test compound for binding the polypeptides. Antibodies can be
25 used to detect the presence of any peptide which shares one or more antigenic determinants with PSEQ or the polypeptides encoded by NSEQ.

In another aspect, the polynucleotides and polypeptides of the present invention can be employed for treatment or the monitoring of therapeutic treatments for cancer. The polynucleotides of NSEQ or those encoding PSEQ, or any fragment or complement thereof,
30 may be used for therapeutic purposes. In one aspect, the complement of the polynucleotides of NSEQ or those encoding PSEQ may be used in situations in which it would be desirable to

block the transcription or translation of the mRNA.

Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. Methods which are well known to those skilled in the art can be used to construct vectors to express nucleic acid sequences complementary to the polynucleotides encoding PSEQ. (See, e.g., Sambrook, supra; and Ausubel, supra.)

Genes having polynucleotide sequences of NSEQ or those encoding PSEQ can be turned off by transforming a cell or tissue with expression vectors which express high levels of a polynucleotide, or fragment thereof, encoding PSEQ. Such constructs may be used to introduce untranslatable sense or antisense sequences into a cell. Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, are preferred. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee et al. (1994) In: Huber and Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. Alternatively, nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases may be included.

Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g.,

Goldman et al. (1997) Nature Biotechnology 15:462-466.)

Further, an antagonist or antibody of a polypeptide of PSEQ or encoded by NSEQ may be administered to a subject to treat or prevent a cancer associated with increased expression or activity of PSEQ. An antibody which specifically binds the polypeptide may be used
5 directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissue which express the the polypeptide.

Antibodies to PSEQ or polypeptides encoded by NSEQ may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments
10 produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are especially preferred for therapeutic use. Monoclonal antibodies to PSEQ may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. In
15 addition, techniques developed for the production of chimeric antibodies can be used. (See, for example, Meyers, supra.) Alternatively, techniques described for the production of single chain antibodies may be employed. Antibody fragments which contain specific binding sites for PSEQ or the polypeptide sequences encoded by NSEQ may also be generated.

Various immunoassays may be used for screening to identify antibodies having the
20 desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art.

Yet further, an agonist of a polypeptide of PSEQ or that encoded by NSEQ may be administered to a subject to treat or prevent a cancer associated with decreased expression or
25 activity of the polypeptide.

An additional aspect of the invention relates to the administration of a pharmaceutical or sterile composition, in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of polypeptides of PSEQ or those encoded by NSEQ, antibodies to the polypeptides, and
30 mimetics, agonists, antagonists, or inhibitors of the polypeptides. The compositions may be administered alone or in combination with at least one other agent, such as a stabilizing

compound, which may be administered in any sterile, biocompatible pharmaceutical carrier including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs, or hormones.

5 The pharmaceutical compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

10 In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA).

15 For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells or in animal models such as mice, rats, rabbits, dogs, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

20 A therapeutically effective dose refers to that amount of active ingredient, for example, polypeptides of PSEQ or those encoded by NSEQ, or fragments thereof, antibodies of the polypeptides, and agonists, antagonists or inhibitors of the polypeptides, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by
25 calculating the ED_{50} (the dose therapeutically effective in 50% of the population) or LD_{50} (the dose lethal to 50% of the population) statistics.

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

30

EXAMPLES

It is understood that this invention is not limited to the particular methodology,

protocols, and reagents described, as these may vary. It is also understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims. The examples below are provide to illustrate the subject invention and are
5 not included for the purpose of limiting the invention.

I cDNA Library Construction

The cDNA library, THYMFET02, was selected to demonstrate the construction of the cDNA libraries from which novel matrix remodeling genes were derived. The THYMFET02 cDNA library was constructed from microscopically normal thymus tissue obtained from a
10 Caucasian female fetus who died at 17 weeks gestation from anencephaly. Serology was negative; family history included tobacco abuse and gastritis.

The frozen tissue was homogenized and lysed in TRIZOL reagent (1 gm tissue/10 ml TRIZOL; Life Technologies, Rockville MD), a monoplastic solution of phenol and guanidine isothiocyanate, using a POLYTRON homogenizer (PT-3000; Brinkmann Instruments,
15 Westbury NY). After a brief incubation on ice, chloroform was added (1:5 v/v), and the lysate was centrifuged. The upper chloroform layer was removed, and the RNA was precipitated with isopropanol, resuspended in DEPC-treated water, and treated with DNase for 25 min at 37°C. The mRNA was reextracted once with acid phenol-chloroform pH 4.7 and precipitated using 0.3 M sodium acetate and 2.5 volumes ethanol. The mRNA was isolated using the
20 OLIGOTEX kit (Qiagen, Chatsworth CA) and used to construct the cDNA library.

The mRNA was handled according to the recommended protocols in the SUPERScript Plasmid system (Life Technologies). The cDNAs were fractionated on a SEPHAROSE CL4B column (Amersham Pharmacia Biotech, Piscataway NJ), and those cDNAs exceeding 400 bp were ligated into pINCY 1 plasmid (Incyte Pharmaceuticals, Palo
25 Alto CA). The plasmid was subsequently transformed into DH5a competent cells (Life Technologies).

II Isolation and Sequencing of cDNA Clones

Plasmid DNA was released from the cells and purified using the REAL Prep 96 Plasmid kit (Qiagen). This kit enabled the simultaneous purification of 96 samples in a 96-
30 well block using multi-channel reagent dispensers. The recommended protocol was employed except for the following changes: 1) the bacteria were cultured in 1 ml of sterile Terrific Broth

(Life Technologies) with carbenicillin at 25 mg/L and glycerol at 0.4%; 2) after inoculation, the cultures were incubated for 19 hours and at the end of incubation, the cells were lysed with 0.3 ml of lysis buffer; and 3) following isopropanol precipitation, the plasmid DNA pellet was resuspended in 0.1 ml of distilled water. After the last step in the protocol, samples were
5 transferred to a 96-well block for storage at 4°C.

The cDNAs were prepared using a MICROLAB 2200 (Hamilton, Reno NV) in combination with DNA ENGINE thermal cyclers (PTC200; MJ Research, Watertown MA) and sequenced by the method of Sanger et al. (1975, J Mol Biol 94:441f) using ABI PRISM 377 DNA sequencing systems.

10 III Selection, Assembly, and Characterization of Sequences

The sequences used for coexpression analysis were assembled from EST sequences, 5' and 3' longread sequences, and full length coding sequences. Selected assembled sequences were expressed in at least three cDNA libraries.

The assembly process is described as follows. EST sequence chromatograms were
15 processed and verified. Quality scores were obtained using PHRED (Ewing et al. (1998) Genome Res 8:175-185; Ewing and Green (1998) Genome Res 8:186-194). Then the edited sequences were loaded into a relational database management system (RDBMS). The EST sequences were clustered into an initial set of bins using BLAST with a product score of 50. All clusters of two or more sequences were created as bins. The overlapping sequences
20 represented in a bin correspond to the sequence of a transcribed gene.

Assembly of the component sequences within each bin was performed using a modification of PHRAP, a publicly available program for assembling DNA fragments (Phil Green, University of Washington, Seattle WA). Bins that showed 82% identity from a local pair-wise alignment between any of the consensus sequences were merged.

25 Bins were annotated by screening the consensus sequence in each bin against public databases, such as GBpri and GenPept from NCBI. The annotation process involved a FASTn screen against the GBpri database in GenBank. Those hits with a percent identity of greater than or equal to 70% and an alignment length of greater than or equal to 100 base pairs were recorded as homolog hits. The residual unannotated sequences were screened by FASTx
30 against GenPept. Those hits with an E value of less than or equal to 10^{-8} are recorded as homolog hits.

Sequences were then reclustered using BLASTn and Cross-Match, a program for rapid protein and nucleic acid sequence comparison and database search (Green, supra), sequentially. Any BLAST alignment between a sequence and a consensus sequence with a score greater than 150 was realigned using cross-match. The sequence was added to the bin
5 whose consensus sequence gave the highest Smith-Waterman score amongst local alignments with at least 82% identity. Non-matching sequences created new bins. The assembly and consensus generation processes were performed for the new bins.

IV Coexpression Analyses of Known Matrix-remodeling Genes

Twenty one known matrix-remodeling genes were selected to identify novel genes that
10 are closely associated with matrix remodeling. The known genes were osteonectin (BM-40), chondroitin/dermatan sulfate proteoglycans (C/DSPG), collagen I, II, III, and IV (coll-I, coll-II, and coll-III), connective tissue growth factor (CTGF), fibrillin, fibronectins, fibronectin receptor (fibr-r), fibulin 1, heparan sulfate proteoglycans (HSPG), extracellular matrix protein (hevin), insulin-like growth factor 1 (IGF 1), insulin-like growth factor binding protein
15 (IGFBP), laminin, lumican, matrix Gla protein (MGP), matrix metalloproteases (MMP), and tissue inhibitors of matrix metalloproteinase 1, 2, and 3 (TIMP 1, 2, and 3). The protein products of the known matrix-remodeling genes may be categorized as follows.

1. Extracellular matrix component protein. These proteins include collagens, proteoglycans, fibrillin, fibronectin, fibulin, and laminin that constitute the major structures of
20 the extracellular matrix.
2. Matrix proteases and matrix protease inhibitors. These proteins include matrix metalloproteases (MMPs) such as the collagenases, and MMP inhibitors such as the tissue-inhibitors of matrix metalloproteases (TIMPs).
3. Regulatory proteins that control expression of matrix-remodeling genes. Such
25 regulatory proteins include connective tissue growth factor, insulin-like growth factor, osteonectin (BM-40), and the receptors for and inhibitors of these proteins.

The known matrix-remodeling genes that we examined in this analysis, and brief descriptions of their functions, are listed in Table 4. Detailed descriptions of their roles in matrix remodeling may be found in the cited articles and reviews.

30

Table 4. Known Matrix-remodeling Genes.

Gene	Description & References
BM-40	<p>Alternate names: SPARC, osteonectin</p> <p>Regulates connective tissue remodeling, wound healing, angiogenesis</p> <p>Induces matrix metalloprotease synthesis (collagenase & gelatinase)</p> <p>Regulates cell movement and proliferation</p> <p>Expression increased in neoplastic melanoma, fibrosis, angiogenesis.</p> <p>(Kamihagi et al. (1994) <i>Biochem Biophys Res Commun</i> 200:423-8; Lane et al. (1994) <i>J Cell Biol</i> 125:929-43; Inagaki et al. (1996) <i>Life Sci</i> 58:927-34; Ledda et al. (1997) <i>J Invest Dermatol</i> 108:210-4; Shankavaram et al. (1997) <i>J Cell Physiol</i> 173:327-34.)</p>
C/DSPG	<p>Chondroitin/dermatan sulfate proteoglycans</p> <p>Major extracellular matrix proteoglycan</p> <p>Regulate cell proliferation, attachment and migration.</p> <p>Darnell et al. (1990) <i>Molecular Cell Biology</i>, Scientific American Press, New York NY; Toole (1991) In: <i>Cell Biology of Extracellular Matrix</i>, Plenum, New York NY, pp. 305-341; Beck et al. (1993) <i>Biochem Biophys Res Commun</i> 190:616-23)</p>
Collagens	<p>Family of fibrous structural proteins (collagen I, II, III, IV, etc.)</p> <p>Most abundant structural component of the extracellular matrix</p> <p>Secreted as procollagen; converted to collagen by MMPs</p> <p>(Alexander and Werb (1991) In: <i>Cell Biology of Extracellular Matrix</i>, pp. 255-302; Adams (1993) In: <i>Extracellular Matrix</i>, Marcel Dekker, New York, NY pp. 91-119; Schuppan et al. (1993) In: <i>Extracellular Matrix</i>, pp. 201-254.)</p>
CTGF	<p>Connective tissue growth factor</p> <p>Mediates induction of matrix synthesis and fibrosis</p> <p>(Grotendorst (1997) <i>Cytokine Growth Factor Rev</i> 8:171-9; Oemar and Luscher (1997) <i>Arterioscler Thromb Vasc Biol</i> 17:1483-9; Ito et al. (1998) <i>Kidney Int</i> 53:853-61.)</p>
fibrillin	<p>Major component of extracellular microfibrills (matrix elastic network)</p> <p>Present in connective tissue throughout the body</p> <p>(Kielty and Shuttleworth (1995) <i>Int J Biochem Cell Biol</i> 27:747-60; Haynes et al. (1997) <i>Br J Dermatol</i> 137:17-23; Hayward and Brock (1997) <i>Hum Mutat</i> 10:415-23.)</p>
fibronectins	<p>Family of extracellular matrix glycoproteins</p> <p>Anchor cells to the matrix</p> <p>Bind matrix proteins to cell surface receptors</p>
fibr-r	<p>Fibronectin receptor</p> <p>Fibronectin receptors regulate cell adhesion & migration</p> <p>(Darnell et al. (1990) <i>Molecular Cell Biology</i>, Scientific American Press, New York NY; Ruoslahti (1991) <i>Cell Biology of Extracellular Matrix</i>, pp. 343-363; Yamada (1991) <i>Cell Biology of Extracellular Matrix</i>, pp. 111-146.)</p>
fibulin 1	<p>Fibronectin-binding extracellular matrix protein</p>

		Mediates platelet adhesion via a bridge of fibrinogen Cleaved by matrix metalloproteinases Inhibits breast and ovarian cancer cell motility (Argaves et al. (1990) J Cell Biol 111:3155-64; Sasaki et al. (1996) Eur J Biochem 240:427-34; Hayashido et al. (1998) Int J Cancer 75:654-8.)
5	HSPG	Heparan sulfate proteoglycans Extracellular matrix proteoglycan found on cell surface of many cell types Regulate cell interactions with the extracellular matrix Bind to collagens and fibronectin in the matrix
10		Regulate cell proliferation, attachment and migration (Darnell et al. (1990) ; Toole (1991) In: <u>Cell Biology of Extracellular Matrix</u> , pp. 305-341; Schuppan et al. (1993) In: <u>Extracellular Matrix</u> , pp. 201-254.)
	hevin	Extracellular matrix protein Homolog to BM-40
15		Regulates cell adhesion and migration Downregulated in metastatic prostate cancer, lung cancer (Girard and Springer (1996) J Biol Chem 271:4511-7; Bendik et al. Cancer Res 58:232-6.)
	IGF 1	Insulin-like growth factor Regulates matrix homeostasis and remodeling
20		Regulates aggregation, growth and survival of cancer cells (Aston et al. (1995) Am J Respir Crit Care Med 151:1597-603; Bitar and Labbad (1996) J Surg Res 61:113-9; Guvakova and Surmacz (1997) Exp Cell Res 231:149-62; Sunic et al. (1998) Endocrinology 139:2356-62.)
	IGFBP	Insulin-like growth factor binding protein Regulates IGF-1 bioavailability (binds IGF-1 more strongly than the receptor)
25		Degraded by matrix metalloproteases (Kiefer et al. (1991) Biochem Biophys Res Commun 176:219-25; Fowlkes et al. (1995) Prog Growth Factor Res 6:255-63; Parker et al. (1996) J Biol Chem 271:13523-9.)
	laminin	Major protein in basal lamina, with collagen, HSPG, and entactin Anchors cells to the matrix by binding collagen, HSPG and heparin
30		Laminins and collagens are the main targets of MMPs Regulates cell attachment, migration, growth, and differentiation (Yamada et al. (1993) In: <u>Extracellular Matrix</u> , pp. 49-66; Giannelli et al. (1997) Science 277:225-8; Quaranta and Plopper (1997) Kidney Int 51: 1441-6; Soini et al. (1997) Hum Pathol 28:220-6.)
	lumican	Extracellular proteoglycan Organizes collagen fibrils in extracellular matrix
35		(Dourado et al. (1996) Osteoarthritis Cartilage 4:187-96; Scott (1996) Biochemistry 35:8795-9; Cs-Szabo et al. (1997) Arthritis Rheum 40:1037-45.)
	MGP	Matrix Gla protein Regulates calcification of cartilage

Marker for osteoblast activity.

(Shanahan et al. (1994) J Clin Invest 93:2393-402; Luo et al. (1997) Nature 386:78-81; Martinetti et al. (1997) Tumour Biol 18:197-205)

5	MMP	Family of Matrix Metalloproteases (including collagenases) Cleave procollagen to produce collagen (Alexander and Werb (1991) In: <u>Cell Biology of Extracellular Matrix</u> , pp. 255-302; Adams (1993) In: <u>Extracellular Matrix</u> , pp. 91-119; Schuppan et al. (1993) In: <u>Extracellular Matrix</u> pp. 201-254.)
10	TIMP 1, 2, 3	Tissue inhibitors of matrix metalloproteinases Bind and inactivate matrix proteases (Schuppan et al. (1993) In: <u>Extracellular Matrix</u> , pp. 201-254; Zvibel and Kraft (1993) In: <u>Extracellular Matrix</u> , pp. 559-580.)

The coexpression of the 21 known genes with each other is shown in Table 5. The entries in Table 5 are the negative log of the p-value ($-\log p$) for the coexpression of the two genes. As shown, the method successfully identified the strong association of the known genes among themselves, indicating that the coexpression analysis method of the present invention was effective in identifying genes that are closely associated with matrix remodeling.

Table 5. Coexpression of 21 known matrix-remodeling genes. ($-\log p$)

	laminin	fibrillin	lumican	coll IV	TIMP-1	IGFBP	coll VI	TIMP-3	CTGF	hevin	fibulin	BM-40	TIMP-2	HSPG	fibronec	MGP	C/DSP	fibr-r	coll-I	coll-III	MMP	
	laminin	7	9	21	9	15	8	4	5	7	14	10	7	11	9	19	11	7	16	10	13	
	fibrillin	7		13	8	6	7	14	11	4	7	12	7	8	4	8	6	13	6	11	12	11
	lumican	9	13		24	17	16	28	17	17	14	15	22	10	8	12	25	33	14	32	34	17
25	coll IV	21	8	24		17	22	22	13	11	14	28	25	12	22	16	27	26	12	34	25	26
	TIMP-1	9	6	17	17		20	15	11	11	6	10	21	15	9	16	20	13	8	14	20	19
	IGFBP	15	7	16	22	20		20	18	16	11	14	18	14	19	21	25	23	10	27	23	20
	coll VI	8	14	28	22	15	20		13	17	19	16	20	11	11	19	19	28	12	31	36	27
	TIMP-3	4	11	17	13	11	18	13		13	18	20	22	14	9	10	18	25	12	12	13	9
30	CTGF	5	4	17	11	11	16	17	13		8	10	18	7	7	19	22	12	12	18	13	11
	hevin	7	7	14	14	6	11	19	18	8		15	18	13	8	8	23	27	10	14	11	8
	fibulin	14	12	15	28	10	14	16	20	10	15		19	9	11	8	19	20	6	17	20	18
	BM-40	10	7	22	25	21	18	20	22	18	18	19		14	11	24	21	24	16	25	32	19
	TIMP-2	7	8	10	12	15	14	11	14	7	13	9	14		7	12	8	16	11	13	13	13
35	HSPG	11	4	8	22	9	19	11	9	7	8	11	11	7		8	14	10	6	11	10	10
	fibronec	9	8	12	16	16	21	19	10	19	8	8	24	12	8		14	14	11	24	21	15
	MGP	19	6	25	27	20	25	19	18	22	23	19	21	8	14	14		32	14	25	20	13
	C/DSPG	11	13	33	26	13	23	28	25	12	27	20	24	16	10	14	32		14	27	28	14
	fibr-r	7	6	14	12	8	10	12	12	12	10	16	16	11	6	11	14	14		14	13	6
40	coll-I	16	11	32	34	14	27	31	12	18	14	17	25	13	11	24	25	27	14		42	21
	coll-III	10	12	34	25	20	23	36	13	13	11	20	32	13	10	21	20	28	13	42		23
	MMP	13	11	17	26	19	20	27	9	11	8	18	19	13	10	15	13	14	6	21	23	

45 V Novel Genes Associated with Matrix Remodeling

Using coexpression analysis, we have identified 20 novel genes that show strong association with known matrix remodeling genes from a total of 41,419 assembled gene

sequences. The degree of association was measured by probability values and has a cutoff of p value less than 0.00001. This was followed by annotation and literature searches to insure that the genes that passed the probability test have strong association with known matrix-remodeling genes. This process was reiterated so that the initial 41,419 genes were reduced to
 5 the final 20 matrix-remodeling genes. Details of the coexpression patterns for the 20 novel matrix-remodeling genes are presented in Table 6.

Each of the 20 novel genes is coexpressed with at least two of the 21 known genes with a p-value of less than 10^{-7} . The coexpression results are shown in Table 6.

The novel genes identified are listed in the table by their Incyte clone numbers (Clone), and
 10 the known genes their abbreviated names (Gene) as shown in Example IV.

Table 6. Coexpression of 20 novel genes with known matrix-remodeling genes. (- log p)

15 Gene	laminin	fibrillin	lumican	coll IV	TIMP-1	IGFBP	coll VI	TIMP-3	CTGF	hevin	fibulin	BM-40	TIMP-2	HSPG	fibronectin	MGP	C/DSP	fibr-r	coll-I	coll-III	MMP
606132	8	7	2	6	4	7	7	2	4	4	4	3	3	4	4	3	2	2	5	3	10
627722	3	4	1	1	3	3	2	5	3	6	3	4	3	2	6	5	3	3	2	3	4
639644	6	7	11	10	3	4	7	3	14	6	6	9	6	2	9	8	5	6	9	7	6
1362659	6	5	6	7	6	9	10	9	8	8	7	6	8	6	7	9	9	7	10	5	5
20 1446685	6	6	11	13	4	7	8	5	7	5	10	9	5	9	5	9	8	6	8	10	7
1556751	3	7	7	8	8	9	9	8	7	6	5	5	7	8	4	10	11	3	7	6	8
1656953	6	8	6	2	5	7	8	5	6	9	3	7	4	3	4	10	8	7	4	4	5
1662318	9	3	6	10	7	9	5	5	8	8	6	8	5	9	6	8	6	4	7	7	9
1996726	3	4	7	7	6	5	8	3	10	2	2	3	2	2	9	3	6	6	8	11	6
25 2137155	3	2	6	3	4	2	2	4	6	4	2	9	4	2	8	4	4	4	5	2	5
2268890	9	13	7	9	8	11	8	9	5	5	8	7	8	5	8	8	11	3	11	7	11
2305981	3	2	4	6	3	4	3	5	5	6	7	5	2	2	2	7	6	4	3	2	2
2457612	3	3	3	5	2	4	4	2	8	4	5	5	2	2	7	8	6	6	5	4	8
2814981	6	3	5	7	4	6	7	2	2	5	5	5	3	6	5	4	6	1	6	4	7
30 3089150	4	6	11	8	5	10	13	9	14	10	11	10	7	6	8	11	16	11	9	7	5
3206667	8	5	10	9	7	5	6	4	9	4	7	8	4	4	7	13	12	4	8	8	6
3284695	7	6	7	14	8	7	6	14	8	18	12	9	10	8	6	18	10	5	13	6	6
3481610	3	2	4	4	3	6	4	6	6	7	4	5	1	5	5	7	5	3	3	2	2
3722004	6	4	8	10	13	9	7	13	8	9	11	12	11	5	10	9	12	3	7	7	6
35 3948614	11	8	6	17	8	13	12	5	5	11	12	7	11	13	4	7	7	4	14	11	10

VI Novel Genes Associated with Matrix Remodeling

The 20 novel genes were identified from the data shown in Table 6 to be associated with matrix remodeling.

40 The nucleotide sequences comprising the consensus sequences of SEQ ID NOs:1-20 of the present invention were first identified from Incyte Clones 606132, 627722, 639644, 1362659, 1446685, 1556751, 1656953, 1662318, 1996726, 2137155, 2268890, 2305981, 2457612, 2814981, 3089150, 3206667, 3284695, 3481610, 3722004, and 3948614, respectively, and assembled according to Example III. BLAST and other motif searches were performed for SEQ ID NOs:1-20 according to
 45 Example VII. The sequences of SEQ ID NOs:1-20 were translated and sequence identity was sought with known sequences. Polypeptide sequences comprising the consensus sequences of SEQ ID NO:21, SEQ ID NO:22, and SEQ ID NO:23 of the present invention were encoded by SEQ ID NO:2,

SEQ ID NO:6, and SEQ ID NO:11, respectively. SEQ ID NOs:21-23 were analyzed using BLAST and other motif search tools as disclosed in Example VII.

SEQ ID NO:3 is 2987 residues in length and shows about 59% sequence identity from about nucleotide 2117 to about nucleotide 2914 with the cDNA encoding regulatory subunit of a human cAMP-dependent protein kinase, RIIBeta (WO 88/03164). SEQ ID NO:8 is 3017 nucleotides in length and shows about 70% to about 74% sequence identity from about nucleotide 1 to about nucleotide 1260 and about nucleotide 1925 to about nucleotide 1985 with human Hpast mRNA (g2529706), a gene associated with multiple endocrine neoplasia type 1. SEQ ID NO:9 is 1735 nucleotides in length and shows about 25% sequence identity from about nucleotide 5 to about nucleotide 1534 with a human neuronal cell adhesion molecule (WO 96/04396) important in the development of nervous system by promoting cell-cell adhesion. SEQ ID NO:14 is 2040 nucleotides in length and shows about 60% to 70% sequence identity from about nucleotide 1 to about nucleotide 1023 with a human mRNA for a serine protease (g1621243) specific for insulin-like growth factor-binding proteins. The amino acid sequence encoded by SEQ ID NO:14 from about nucleotide 3 to about nucleotide 1043 shows about 61% sequence identity with an osteoblast-like cell-derived protein (J09107980) useful for treatment and prevention of various diseases and as contraceptive. SEQ ID NO:15 is 2121 nucleotides in length and shows 60-80% sequence identity with a mouse gene, ADAMT-1 (g2809056), a member of the ADAM (the disintegrin and metalloproteinase) family. ADAMT-1 has been shown to contain the thrombospondin (TSP) type I motif; expression of ADAMT-1 is closely associated with inflammatory processes (Kuno et al (1997) Genomics 46:466-471). SEQ ID NO:16 is 2900 nucleotides in length and shows about 70% sequence identity with a mouse homeobox (Pmx) mRNA (g460124). Homeobox genes are expressed in very specific temporal and spatial pattern and function as transcriptional regulators of developmental processes. (Kern et al. (1994) Genomics 19:334-340).

SEQ ID NO:21 is 551 amino acid residues long and shows about 37% sequence identity from about amino acid residue 10 to about amino acid residue 278 with PALM (g3219602), a human paralemin that is membrane-bound and expressed abundantly in brain and at intermediate levels in the kidney and in endocrine cells. In addition, the sequence encompassing residues 418 to 434 of SEQ ID NO:21 resembles one of the structural fingerprint regions of a seven trans-membrane receptor, LCR1, that is isolated from the human brain (Rimland et al. (1991) Mol Pharmacol 40:869-875). SEQ ID NO:21 also has one potential amidation site at L546; three potential N-glycosylation sites at N223, N229, and N408; one potential cAMP- and cGMP-dependent protein kinase phosphorylation site at S486; fifteen potential casein kinase II phosphorylation sites at S57, S100, T101, T116, S135, S253, T349, S370, T387, S426, T434, S489, S505, S520, and T526; one potential N-myristoylation site at G54; and nine potential protein kinase C phosphorylation sites at T15, S25, S57, S100, S123, S247,

S364, S370, and S505. SEQ ID NO:22 is 99 amino acid residues in length. The sequence of SEQ ID NO:22 from about amino acid residue 71 to about amino acid residue 81 resembles one of the fingerprint regions of the RH1 and RH2 opsins, a family of G protein coupled receptors that mediate vision (Zuker et al. (1985) Cell 40:851-858; Cowman et al. (1986) Cell 44:705-710). SEQ ID NO:22 also has one potential N-myristoylation site at G24, and two potential protein kinase C phosphorylation sites at S13 and S89. SEQ ID NO:23 is 493 amino acid residues in length and shows about 44% sequence identity from about amino acid residue 277 to about amino acid residue 487 with an angiopoietin-like factor from the human cornea, CDT6 (g2765527). Angiopoietin 1 and angiopoietin 2 function as a natural ligand and a natural inhibitor, respectively, for TIE2, a receptor critical in angiogenesis during embryonic development, tumor growth, and tumor metastasis. The sequences encompassing amino acid residues 305 to 343, 346 to 355, 365 to 402, 411 to 424, and 428 to 458 of SEQ ID NO:23 resemble the carboxy-terminal domain signatures of fibrinogen beta and gamma chains from BLOCKS analysis. SEQ ID NO:23 also exhibits one potential signal peptide region encompassing amino acid residues M1 to G22 when analyzed using a HMM-based signal peptide analysis tool. In addition, SEQ ID NO:23 shows two potential N-glycosylation sites at N164 and N192; one potential cAMP- and cGMP-dependent protein kinase phosphorylation sites at S127, six potential casein kinase II phosphorylation sites at S34, S209, T238, S266, T368, and T417; four potential N-myristoylation sites at G12, G18, G22, and G29; eight potential protein kinase C phosphorylation sites at S34, S209, T268, T299, T335, S373, S383, and S477; and three potential tyrosine kinase phosphorylation sites at Y183, Y392, and Y467.

VII Homology Searching for Matrix-Remodeling Genes and the Proteins Encoded by the Genes

Polynucleotide sequences, SEQ ID NOs:1-20, and polypeptide sequences, SEQ ID NOs: 21-23, were queried against databases derived from sources such as GenBank and SwissProt. These databases, which contain previously identified and annotated sequences, were searched for regions of similarity using Basic Local Alignment Search Tool (BLAST; Altschul (1990) supra) and Smith-Waterman alignment (Smith et al. (1992) Protein Engineering 5:35-51). BLAST searched for matches and reported only those that satisfied the probability thresholds of 10^{-25} or less for nucleotide sequences and 10^{-8} or less for polypeptide sequences.

The polypeptide sequences were also analyzed for known motif patterns using MOTIFS, SPSCAN, BLIMPS, and Hidden Markov Model (HMM)-based protocols. MOTIFS (Genetics Computer Group, Madison WI) searches polypeptide sequences for patterns that match those defined in the Prosite Dictionary of Protein Sites and Patterns (Bairoch et al. supra), and displays the patterns found and their corresponding literature abstracts. SPSCAN (Genetics Computer Group) searches for

potential signal peptide sequences using a weighted matrix method (Nielsen et al. (1997) Prot Eng 10:1-6). Hits with a score of 5 or greater were considered. BLIMPS uses a weighted matrix analysis algorithm to search for sequence similarity between the polypeptide sequences and those contained in BLOCKS, a database consisting of short amino acid segments, or blocks, of 3-60 amino acids in length, compiled from the PROSITE database (Henikoff et al. supra; Bairoch et al. supra), and those in PRINTS, a protein fingerprint database based on non-redundant sequences obtained from sources such as SwissProt, GenBank, PIR, and NRL-3D (Attwood et al. (1997) J Chem Inf Comput Sci 37:417-424). For the purposes of the present invention, the BLIMPS searches reported matches with a cutoff score of 1000 or greater and a cutoff probability value of 1.0×10^{-3} . HMM-based protocols were based on a probabilistic approach and searched for consensus primary structures of gene families in the protein sequences (Eddy, supra; Sonnhammer, supra). More than 500 known protein families with cutoff scores ranging from 10 to 50 bits were selected for use in this invention.

VIII Labeling and Use of Individual Hybridization Probes

Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250 μ Ci of [γ - 32 P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (NEN Life Science Products, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine resin column (Amersham Pharmacia Biotech). An aliquot containing 10^7 counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (NEN Life Science Products).

The DNA from each digest is fractionated on a 0.7 percent agarose gel and transferred to nylon membranes (NYTRAN PLUS, Schleicher & Schuell, Durham NH). Hybridization is carried out under the following conditions: 5x SCC/0.1% SDS at 60° C for about 6 hours, subsequent washes are performed at higher stringency with buffers, such as 1x SCC/0.1% SDS at 45° C, then 0.1xSCC. After XOMAT AR film (Eastman Kodak, Rochester NY) is exposed to the blots for several hours, hybridization patterns are compared.

IX Production of Specific Antibodies

SEQ ID NO:20, 21, or 23 substantially purified using polyacrylamide gel electrophoresis (Harrington (1990) Methods Enzymol 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the amino acid sequence is analyzed using LASERGENE software (DNASTAR, Madison WI) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for

selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. Typically, oligopeptides 15 residues in length are synthesized using an ABI 431A peptide synthesizer (PE Biosystems) using Fmoc-chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester to increase

5 immunogenicity. Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide activity by, for example, binding the peptide to plastic, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-

iodinated goat anti-rabbit IgG.

10

What is claimed is:

1. A substantially purified polynucleotide comprising a gene that is coexpressed with one or
5 more known matrix-remodeling genes in a plurality of biological samples, wherein each known
matrix-remodeling gene is selected from the group consisting of osteonectin, chondroitin/dermatan
sulfate proteoglycans, collagen I, II, II, and IV, connective tissue growth factor, fibrillin, fibronectins,
fibronectin receptor, fibulin 1, heparan sulfate proteoglycan, extracellular matrix protein, insulin-like
growth factor 1, insulin-like growth factor binding protein, laminin, lumican, matrix Gla protein,
10 matrix metalloproteases, and tissue inhibitors of matrix metalloproteinase 1, 2, and 3.

2. The polynucleotide of claim 1, comprising a polynucleotide sequence selected from the
group consisting of:

- (a) a polynucleotide sequence selected from the group consisting of SEQ ID NOs:1- 20;
- (b) a polynucleotide sequence which encodes the polypeptide sequence of SEQ ID NO: 21,
15 22, or 23;
- (c) a polynucleotide sequence having at least 70% identity to the polynucleotide sequence of
(a) or (b);
- (d) a polynucleotide sequence comprising at least 18 sequential nucleotides of the
polynucleotide sequence of (a), (b), or (c);
- 20 (e) a polynucleotide which hybridizes under stringent conditions to the polynucleotide of (a),
(b), (c), or (d); and
- (f) a polynucleotide sequence which is complementary to the polynucleotide sequence of (a),
(b), (c), (d), or (e).

3. A substantially purified polypeptide comprising the gene product of a gene that is
25 coexpressed with one or more known matrix-remodeling genes in a plurality of biological samples,
wherein each known matrix-remodeling gene is selected from the group consisting of osteonectin,
chondroitin/dermatan sulfate proteoglycans, collagen I, II, II, and IV, connective tissue growth factor,
fibrillin, fibronectins, fibronectin receptor, fibulin 1, heparan sulfate proteoglycans, extracellular
matrix protein, insulin-like growth factor 1, insulin-like growth factor binding protein, laminin,
30 lumican, matrix Gla protein, matrix metalloproteases, and tissue inhibitors of matrix metalloproteinase
1, 2, and 3.

4. The polypeptide of claim 3, comprising a polypeptide sequence selected from the group
consisting of:

- (a) the polypeptide sequence of SEQ ID NO:21, 22, or 23;
- 35 (b) a polypeptide sequence having at least 85% identity to the polypeptide sequence of (a);
and

(c) a polypeptide sequence comprising at least 6 sequential amino acids of the polypeptide sequence of (a) or (b).

5. An expression vector comprising the polynucleotide of claim 2.

6. A host cell comprising the expression vector of claim 5.

5 7. A pharmaceutical composition comprising the polynucleotide of claim 2 or the polypeptide of claim 3 in conjunction with a suitable pharmaceutical carrier.

8. An antibody which specifically binds to the polypeptide of claim 4.

9. A method for diagnosing a disease or condition associated with the altered expression of a gene that is coexpressed with one or more known matrix-remodeling genes, wherein each known
10 matrix-remodeling gene is selected from the group consisting of osteonectin, chondroitin/dermatan sulfate proteoglycans, collagen I, II, II, and IV, connective tissue growth factor, fibrillin, fibronectins, fibronectin receptor, fibulin 1, heparan sulfate proteoglycans, extracellular matrix protein, insulin-like growth factor 1, insulin-like growth factor binding protein, laminin, lumican, matrix Gla protein, matrix metalloproteases, and tissue inhibitors of matrix metalloproteinase 1, 2, and 3, the method
15 comprising the steps of:

(a) providing a sample comprising one of more of said coexpressed genes;

(b) hybridizing the polynucleotide of claim 2(F) to said coexpressed genes under conditions effective to form one or more hybridization complexes; and

(c) detecting the hybridization complexes, wherein the altered level of hybridization
20 complexes compared with the level of hybridization complexes of a nondiseased sample correlates with the presence of the disease or condition.

10. A method for treating or preventing a disease associated with the altered expression of a gene that is coexpressed with one or more known matrix-remodeling genes in a subject in need, wherein each known matrix-remodeling gene is selected from the group consisting of osteonectin,
25 chondroitin/dermatan sulfate proteoglycans, collagen I, II, II, and IV, connective tissue growth factor, fibrillin, fibronectins, fibronectin receptor, fibulin 1, heparan sulfate proteoglycans, extracellular matrix protein, insulin-like growth factor 1, insulin-like growth factor binding protein, laminin, lumican, matrix Gla protein, matrix metalloproteases, and tissue inhibitors of matrix metalloproteinase 1, 2, and 3, the method comprising the step of administering to said subject in need the pharmaceutical
30 composition of claim 7 in an amount effective for treating or preventing said disease.

11. A method for treating or preventing a disease associated with the altered expression of a gene that is coexpressed with one or more known matrix-remodeling genes in a subject in need, wherein each known matrix-remodeling gene is selected from the group consisting of osteonectin, chondroitin/dermatan sulfate proteoglycans, collagen I, II, II, and IV, connective tissue growth factor,

fibrillin, fibronectins, fibronectin receptor, fibulin 1, heparan sulfate proteoglycans, extracellular matrix protein, insulin-like growth factor 1, insulin-like growth factor binding protein, laminin, lumican, matrix Gla protein, matrix metalloproteases, and tissue inhibitors of matrix metalloproteinase 1, 2, and 3, the method comprising the step of administering to said subject in need the antibody of
5 claim 8 in an amount effective for treating or preventing said disease.

12. A method for treating or preventing a disease associated with the altered expression of a gene that is coexpressed with one or more known matrix-remodeling genes in a subject in need, wherein each known matrix-remodeling gene is selected from the group consisting of osteonectin, chondroitin/dermatan sulfate proteoglycans, collagen I, II, III, and IV, connective tissue growth factor,
10 fibrillin, fibronectins, fibronectin receptor, fibulin 1, heparan sulfate proteoglycans, extracellular matrix protein, insulin-like growth factor 1, insulin-like growth factor binding protein, laminin, lumican, matrix Gla protein, matrix metalloproteases, and tissue inhibitors of matrix metalloproteinase 1, 2, and 3, the method comprising the step of administering to said subject in need the polynucleotide sequence of claim 2(F) in an amount effective for treating or preventing said disease.

SEQUENCE LISTING

<110> INCYTE PHARMACEUTICALS, INC.
WALKER, Michael G.
VOLKMUTH, Wayne
KLINGLER, Tod M.

<120> MATRIX-REMODELING GENES

<130> PB-0004 PCT

<140> To Be Assigned

<141> Herewith

<150> 09/169,289

<151> 1998-10-09

<160> 23

<170> PERL Program

<210> 1

<211> 1447

<212> DNA

<213> Homo sapiens

<220>

<221> unsure.

<222> 1380

<223> a or g or c or t, unknown, or other

<220>

<221> misc feature

<223> Incyte ID No.: 606132CB1

<400> 1

```
cctggaacca gaaggagacc tacctgcaca tcatgaagaa cgaggaggag gtggtgatct 60
tggtcgcgca ggtgggcgac cgcagcatca tgcaaagcca gagcctgatg ctggagctgc 120
gagagcagga ccaggtgtgg gtacgcctct acaagggcga acgtgagaac gccatcttca 180
gcgaggagct ggacacctac atcaccttca gtggctacct ggtcaagcac gccaccgagc 240
cctagctggc cggccacctc ctttcctctc gccaccttcc acccctgcgc tgtgctgacc 300
ccaccgcctc ttccccgatc cctggactcc gactccctgg ctttggcatt cagtgagacg 360
ccctgcacac acagaaagcc aaagcgatcg gtgctcccag atcccgcagc ctctggagag 420
agctgacggc agatgaaatc accagggcgg ggcacccgcg agaaccctct gggaccttcc 480
gcggccctct ctgcacacat cctcaagtga ccccgacggg cgagacgcgg gtggcggcag 540
ggcgtcccag ggtgcggcac cgcggctcca gtccttggaa ataattagga aaattctaaa 600
ggtctcaaaa ggagcaaagt aaaccgtgga ggacaaagaa aagggttgtt atttttgtct 660
ttccagccag cctgctggct cccaagagag aggccttttc agttgagact ctgcttaaga 720
gaagatccaa agttaagct ctggggctcag gggaggggccc gggggcagga aactacctct 780
ggcttaattc ttttaagcca cgtaggaact ttcttgaggg ataggtggac cctgacatcc 840
ctgtggcctt gcccaagggc tctgctggtc tttctgagtc acagctgcga ggtgatgggg 900
gctggggccc caggcgctcag ctcccagagg gacagctgag cccctgcct tggctccagg 960
ttggtagaag cagccgaagg gctcctgaca gtggccaggg acccctgggt ccccaggcc 1020
tgcatatgtt tctatgaggg gcagagctcc tggtagatcc atgtgtggct ctgctccacc 1080
cctgtgccac ccagagccc tggggggtgg tctccatgcc tgccaccctg gcatcggttt 1140
tctgtgccgc ctcccacaca aatcagcccc agaaggcccc ggggccttgg cttctgtttt 1200
ttataaaaca cctcaageag cactgcagtc tcccatctcc tcgtgggcta agcatcaccg 1260
cttccacgtg tgttgtgttg gttggcagca aggctgatcc agacccttc tgccccact 1320
gcgctcatcc aggcctctga ccagtagcct gagaggggct tttctaggc ttcagagcan 1380
gggagagctg gacggggtag acagtccgct tgtctgttct aagctctgtg agctcagtct 1440
gagacaa
```

<210> 2
<211> 2481
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No.: 627722CB1

<400> 2
ctagcaagca ggtaaacgag ctttgtacaa acacacacag accaacacat ccgggggatgg 60
ctgtgtgttg ctagagcaga ggctgattaa acactcagtg tggttgctct ctgtgccact 120
cctggaaaat aatgaattgg gtaaggaaca gttaataaga aaatgtgcct tgctaactgt 180
gcacattaca acaaagagct ggcagctcct gaaggaaaag ggcttgtgcc gctgccgttc 240
aaacttgtca gtcaactcat gccagcagcc tcagcgtctg cctccccagc acaccctcat 300
tacatgtgtc tgtctggcct gatctgtgca tctgtctcga gacgctcctg acaagtcggg 360
aattttctta tttctccact ggtgcaaaga gcggatttct ccctgcttct cttctgtcac 420
ccccgctcct ctcccccagg aggtcctctg atttatggta gctttggact tgcttccccg 480
tctgactgtc cttgacttct agaatggaag aagctgagct ggtgaaggga agactccagg 540
ccatcacaga taaaagaaaa atacaggaag aaatctcaca gaagcgtctg aaaatagagg 600
aagacaaact aaagcaccag catttgaaga aaaaggcctt gagggagaaa tggcttctag 660
atggaatcag cagcggaaaa gaacaggaag agatgaagaa gcaaaatcaa caagaccagc 720
accgatcca ggttctagaa caaagtatcc tcaggcttga gaaagagatc caagatcttg 780
aaaaagctga actgcaaatc tcaacgaagg aagaggccat tttaaagaaa ctaaagtcaa 840
ttgagcggac aacagaagac attataagat ctgtgaaagt ggaaagagaa gaaagagcag 900
aagagtcaat tgaggacatc tatgctaata tccctgacct tccaaagtcc tacatacctt 960
ctaggttaag gaaggagata aatgaagaaa aagaagatga tgaacaaaat aggaaagctt 1020
tatatgccat ggaaattaaa gttgaaaaag acttgaagac tggagaaagt acagttctgt 1080
cttcaatacc tctgccatca gatgacttta aaggtacagg aataaaaagt tatgatgatg 1140
ggcaaaagtc agtgtatgca gtaagttcta atcacagtgc agcatacaat ggcaccgatg 1200
gcctggcacc agttgaagta gaggaacttc taagacaagc ctcagagaga aactctaaat 1260
ccccaacaga gtatcatgag cctgtatatg ccaatccctt ttacaggcct acaacccac 1320
agagagaaac ggtgaccctt ggaccaaaact ttcaagaaag gataaagatt aaaactaatg 1380
gactgggtat tgggtgtaaat gaatccatac acaatatggg caatggtctt tcagaggaaa 1440
ggggaaacaa cttcaatcac atcagtccca ttccgccagt gctcatccc cgatcagtga 1500
ttcaacaagc agaagagaag cttcacaccc cgcaaaaag gctaatagact ccttggaag 1560
aatcgaatgt catgcaggac aaagatgcac cctctccaaa gccaaaggct agccccagag 1620
agacaatatt tgggaaatct gaacaccaga attcttcacc cacttgtcag gaggacgagg 1680
aagatgtcag atataatatc gttcattccc tgcctccaga cataaatgat acagaaccgg 1740
tgacaatgat tttcatgggg tatcagcagg cagaagacag tgaagaagat aagaagtttc 1800
tgacaggata tgatgggatc atccatgctg agctggttgt gattgatgat gaggaggagg 1860
aggatgaagg agaagcagag aaaccgtcct accaccccat agtccccat agtcagggtg 1920
accagccagc caaaccaaca ccacttccta gaaaaagatc agaagctagt cctcatgaaa 1980
acacaaatca taaatcccc cacaataatt ccatatctct gaaagagcaa gaagaaagct 2040
taggcagccc tgtccaccat tccccatttg atgctcagac aactggagat gggactgagg 2100
atccatcctt aacagcttta aggatgagaa tggcaaaagt gggaaaaaag gtgatctaag 2160
agttgtacca cctatataaa catcctttga agaagaaact aagaagcatt tgcaaatttc 2220
tcttctggat attttgttta ttttttctga agtccaaaaa attatcatta cagtgtacca 2280
tattaagcca tgtgaataag tagtagtcat tatttgtgaa aaattcccaa aaagctgggg 2340
aaaacaaatg tgtaactttt ccagttactt gacacgattc agtgggggaa aaccagcatt 2400
ttttattcta ttgataccaa agcatttcta ataagagctt gttaaattta agaataaagt 2460
tatttaaaat aaaaaaaaaa a 2481

<210> 3
<211> 2987
<212> DNA
<213> Homo sapiens

<220>
<221> unsure

<222> 2955

<223> a or g or c or t, unknown, or other

<220>

<221> misc_feature

<223> Incyte ID No.: 639644CB1

<400> 3

```
agaaaaaaag aaaaaagaaa aaaactaagg cagcagctct taataaataa cacctggagc 60
agaatcggtg aactgctttc acgttggctt ttgcagaagt ggcaatgcat tgaggatata 120
tctggcaagc ttcgaattca caagtgtaaa ggaccagtg acctgctcac agtccggcag 180
agcacgcgga acctctacgc tcgcggcttc catgacaaag acaaagagtg cagttgtagg 240
gagtctggtt accgtgccag cagaagccaa agaaagagtc aacggcaatt cttgagaaac 300
caggggactc caaagtacaa gccagatgtt gtccatactc ggcagacacg ttccttgtcc 360
gtcgaatttg aaggtgaaat atatgacata aatctggaag aagaagaaga attgcaagtg 420
ttgcaaccaa gaaacattgc taagcgtcat gatgaaggcc acaaggggcc aagagatctc 480
caggcttcca gtggtggcaa caggggcagg atgctggcag atagcagcaa cgccgtgggc 540
ccacctacca ctgtccgagt gacacacaag tgttttattc ttcccaatga ctctatccat 600
tgtgagagag aactgtacca atcggccaga gcgtggaagg accataaggc atacattgac 660
aaagagattg aagctctgca agataaaatt aagaatttaa gagaagtga aggacatctg 720
aagagaagga agcctgagga atgtagctgc agtaaacaaa gctattacaa taaagagaaa 780
ggtgtaaaaa agcaagagaa attaaagagc catcttcacc cattcaagga ggctgctcag 840
gaagtagata gcaaactgca acttttcaag gagaacaacc gtaggaggaa gaaggagagg 900
aaggagaaga gacggcagag gaagggggaa gagtgacagc tgccctggcc cacttgcttc 960
acgcatgaca acaaccactg gcagacagcc ccgttctgga acctgggatc tttctgtgct 1020
tgcacgagtt ctaacaataa cacctactgg tgtttgcgta cagttaatga gacgcataat 1080
tttcttttct gtgagtttgc tactggcttt ttggagtatt ttgatatgaa tacagatcct 1140
tatcagctca caaatacagt gcacacggta gaacgaggca ttttgaatca gctacacgta 1200
caactaatgg agctcagaag ctgtcaagga tataagcagt gcaacccaag acctaagaat 1260
cttgatgttg gaaataaaga tggaggaagc tatgacctac acagaggaca gttatgggat 1320
ggatgggaag gttaatcagc cccgtctcac tgcagacatc aactggcaag gcctagagga 1380
gctacacagt gtgaatgaaa acatctatga gtacagacaa aactacagac ttagtctggt 1440
ggactggact aattacttga aggatttaga tagagtattt gcactgctga agagtacta 1500
tgagcaaaat aaaacaaata agactcaaac tgctcaaagt gacgggttct tggttgtctc 1560
tgctgagcac gctgtgtcaa tggagatggc ctctgctgac tcagatgaag acccaaggca 1620
taagggtggg aaaacacctc atttgacctt gccagctgac cttcaaacc ctcatttgaa 1680
ccgaccaaca ttaagtccag agagtaaact tgaatggaat aacgacattc cagaagttaa 1740
tcatttgaat tctgaacact ggagaaaaac cgaaaaatgg acggggcatg aagagactaa 1800
tcattctgaa accgatttca gtggcgatgg catgacagag ctagagctcg ggcccagccc 1860
caggctgcag cccattcgca ggcacccgaa agaacttccc cagtatgggt gtccctggaa 1920
ggacattttt gaagatcaac tatacttccc tgtgcattcc gatggaattt cagttcatca 1980
gatgttcacc atggccaccg cagaacaccg aagtaattcc agcatagcgg ggaagatgtt 2040
gaccaaggtg gagaagaatc acgaaaagga gaagtcacag cacctagaag gcagcgcctc 2100
ctcttcactc tctctgatt agatgaaact gttaccttac cctaaacaca gtatttcttt 2160
ttaacttttt tatttgtaaa ctaataaagg taatcacagc caccaacatt ccaagctacc 2220
ctgggtacct ttgtgcagta gaagctagt agcatgtgag caagcgggtg gcacacggag 2280
actcatcggt ataatttact atctgccaa agtagaaaga aaggctgggg atatttgggt 2340
tggcttgggt ttgatttttt gcttgtttgt ttgttttgta ctaaaacagt attatctttt 2400
gaatatcgta gggacataag tatatacatg ttatccaatc aagatggcta gaatgggtgcc 2460
tttctgagtg tctaaaactt gacacccctg gtaaatcttt caacacactt ccactgcctg 2520
cgtaatgaag ttttgattca tttttaacca ctggaatttt tcaatgccgt cattttcagt 2580
tagatgattt tgcactttga gattaaaatg ccatgtctat ttgattagtc ttattttttt 2640
atttttacag gcttatcagt ctactgttg gctgtcattg tgacaaagtc aaataaaccc 2700
ccaaggacga cacacagtat ggatcacata ttgtttgaca ttaagctttt gccagaaaat 2760
gttgcatgtg ttttacctcg acttgctaaa atcgattagc agaaagcat ggctaataat 2820
gttggtgggt aaaataaata aataagtaaa caaaaaaaa aaaaaaaaaa aaaaaaaaaa 2880
aaaaaaaaaa aaaaaaaaaa aaaaagcaaa aaaagctgcc gccacagtta gatgaagaag 2940
catgaggatc cgagnnggtc gcctctttga gtggtgaggg agtcgcg 2987
```

<210> 4

<211> 2915

<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No.: 1362659CB1

<400> 4
gaggcaagaa ttccggcacga gggacatttt gccaaacttaa acgagaaaaa gaccccccg 60
acccggcaca ctcccccttc ctccagcccc gcttcagcca catgctccag ctgctgcca 120
gtaaagccct gtgccttttt ttccccctgaa tactgcccac agcatccccct tcccatctgc 180
ctctcaggag ttggggactt tgctaggaga ttttttaagt gttccttact gggacaacgt 240
ggagccacgt ttgcaggagc tccatttgta tccctgctgg tgttgacttc tgtgtagggg 300
ccagttcatg tccctgactc tcacctccca ttagataaat gaagcccacc cccctttcta 360
gagtgatgag agtcaagaag aggggatgta tgaacggcca aattcccatg tgagaggaag 420
atgacctgat ccacctagcc ttttcttctg gatctgtcct ccctcacccc tttcacctga 480
gctgtccaca gtaggaaaca taaagaaaca atgtccccta catatcccca tgactacata 540
atccatcatc gtaggaaata ggaaagcaaa tttgattttg gttttgtaaa acgtacatgc 600
ttcaataatt ctttttttgt gtcttaaata ctcatagggg aaaaaaacag ctcacccaag 660
gtgttaggtt tcacatatat attcatcaac tatttttagaa gatttaattc tatcaaatct 720
tgtattacct cagatcattt taaatagcaa gccaaataacg agctttgaag gctattttac 780
cattcctgtt caaaaaagg tctcatgggt cctgacaggt tacccttgag ggcttgtgtc 840
tactttttta aagtcaatgg ttttttttct tgtgttctag tttccataat aggagagaaa 900
atatagaaat atatgcaaaa attatagttt tcttttagatc agaaactgat atttttgggt 960
cagccatatg tattttgttt aaaggattta aaataaagtg ccgtcatgta gccctgtgga 1020
agggagcaca taaccagctg tttggcatga cagggtgactt agtatatttg taattgggtt 1080
taaaaccaat acaccatact ttctttctgc aaacagccat ctttatactt agggaagaaa 1140
aattgttggg ttctagactt ttttaataata aattttgttg atatggaatt aggtaagttt 1200
aagtgtctat gtgcatatgt tttttatata agttttttct attcagtttc actgatccaa 1260
ctggcagtgg gtaaatatgg cataagttta taacactttt ccccaaaatg gtgctttgga 1320
tttgaaaagg gtctgatggg gagaaggaga acgtatcatc ctagcttcct ctcttaataa 1380
acctagaaaa acgggtagta aactgtggat agtcaggaaa acaccagca agggacacag 1440
ctgtcaggaa atgaatcttc cccccaaccc ccaccatgca gatggataga cagaatcttt 1500
cctgactagt cattaggatc aggggcctct gttggatttg tgtttcttga agaataagctg 1560
gcagagtggg ataaaagaca cgaatatctc ctggtctata aggatactct gatttggggg 1620
ttgcattttt catggttttt atttctgtgt cccctggag ttttccatta gtgagttttt 1680
gtgcaaggat cttatttggt atgccttccc tcccctagaa agattttgtg caatatatta 1740
aatggggaca gaattctaaa tggataaaac aatggctggg tctagccctg agtgacagtc 1800
ttaaggctag atccttccca tagtatcatc tgtcctctgg aatgactctc ctgtccctaa 1860
aggggttaag agagagatca cctagaaatc cctctggaca cttgtgggtt ctttaggggt 1920
tgagtttctt ctcccccctg agcttcagag aggagagttg gcatgggtaa atctgaatgg 1980
ttacctcact gctgaaaacc cagaggggag tggcacactc gcttgtgtgg aaaagcctct 2040
aaatgcatec ctctctttct ttctgtcttc ctttgcctta caattgaagc agcccgtggg 2100
accatcacag tatgcagaga ctctctcacc tttcatatct agggaccacc cccgatgcat 2160
tggtgagggg gggcacttat aaatgcctgc tattgttaag ccattccagc ctcttctct 2220
gaatagacca gacgcccttt cacttagttc agtgccagtc cttttgcctt cccaaccctg 2280
ctgttagggc tgctgttccc tttgtctctg attaggagag atggaaggag atgagctccc 2340
ataactgaat tggccttttg ttcatgtttt ctccccatat gtatatatgc catatgtgaa 2400
tatgccatat atatgtgcca acaaacttat ctacgttggt cttttcaaat tagcacgcag 2460
ataggaattt tgagtttctt ctctcttttag taactagtat aacaagcact ggtatttttg 2520
tacaaaaaag aaaaacaaaa gattgactat tgtggtctgc atgacataaa caaacaaatg 2580
gtgatataca agcaacgtat accccagtc agtggtgtgt gccataattt gcaattcagc 2640
ttaacagtgc acccaatcta tatttgcatt ttgatattat ttaagctcta tgtacaagg 2700
tttgcatgta tttatatggg tcttagggaa aaaaaatgct ataaactgca aatctgaaat 2760
tcaaagtgtg tgttccactg agaccagaag aagaagagga gttttaaaag ggataatttg 2820
ttggagccaa taaagctttt tgctgatgaa cagaaaccaa tactgctgtg cactgagaat 2880
aaaaactcat gccacttgt aaaaaaaaaa aaagg 2915

<210> 5

<211> 1826

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No.: 1446685CB1

<400> 5

```
gaaagccgca gcctcagtec cgccgcccgc cgctgcgtcc gccagcgcc agctccgcgt 60
cccagccggc ccgcggcagc ctgcgcccgc ccatggccac ctccccgcag aagtcgcctt 120
ctgtcpccaa gtctcccact cccaagtcgc ccccgctccg caagaaagat gattccttct 180
tggggaaact cggagggacc ctggcccggg ggaagaaagc caaggagggtg tccgagctgc 240
aggaggaggg aatgaacgcc atcaacctgc ccctcagccc aattcccttt gagctggacc 300
ccgaggacac gatgctggag gagaatgagg tgcgaacaat ggtggatcca aactcacgca 360
gtgaccccaa gttcaagaa ctgatgaagg tattaattga ctggattaat gatgtgttgg 420
ttggagaaaag aatcattgtg aaagacctag ctgaagattt gtatgatgga caagtcctgc 480
agaagctttt cgagaaactg gagagtgaga agctaaatgt ggctgaggtc acccagtcag 540
agattgctca gaagcaaaaa ctgcagactg tcctggagaa gatcaatgaa accctgaaac 600
ttcctcccag gagcatcaag tggaatgtgg attctgttca tgccaagagc ctggtggcca 660
tcttacacct gctcgttgct ctgtctcagt atttccgcgc accaattcga ctcccagacc 720
atgtttccat ccaagtgggt gtggtccaga aacgagaagg aatcctccag tctcggcaaa 780
tccaagagga aataactggt aacacagagg ctctttccgg gaggcataaa cgtgatgcct 840
ttgacacctt gttcgacat gccccagaca agctgaatgt ggtgaaaaag aactcatca 900
ctttcgtgaa caagcacctg aataaactga acctggaggc cacagaactg gaaacccagt 960
ttgcagatgg ggtgtacctg gtgctgctca tggggctcct ggagggtac tttgtgcccc 1020
tgcacagctt ctctctgacc ccggacagct ttgaacagaa ggtcttgaat gtctcctttg 1080
cctttgagct catgcaagat ggagggttgg aaaagccaaa accgcggcca gaagacatag 1140
tcaactgtga cctgaaatct acactacgag tgttgtacaa cctcttcacc aagtaccgta 1200
acgtggagtg aggggctgcc ctggggcccac cactgcccac gagttcttgc tgttggcgta 1260
ctggaccctc ctccgaactg ccttaccctg cttattcctg tctcttgac tgtgctctcc 1320
cacaagtcca gctgcaacct agagatagtg gaaactgaaa ttaggaagga aatcatcaat 1380
aactcagtg gctgacccat ccctcccagg cgctggggac caacctagca atgaagggtg 1440
ggaagggtgt tcccttcccg gtgccagggtc cagatttccc tccatgattt gggaaccagg 1500
ttaggcaaaa gagtccccac aagatgaaaa taaagatcct agttaccatt caaaggatgc 1560
taactgtgtg tcaggcccca cactaagtgc tctgctctga tatactcaag gccattaatc 1620
ttcaggactc ccattgacgt aggtgtttca ttcccctttt acagatgagg aaactaaggc 1680
ttggagggtta aatgacttgc cagaagttgg aatttttttc ctctttgaac ataacctctc 1740
ccttctccct aaaggtaacc actattctga gtccaatcat caagggtttg cttttctttt 1800
tagctaagta tgcattcctc aatagt 1826
```

<210> 6

<211> 1439

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No.: 1556751CB1

<400> 6

```
gagtatccct tgtttaatca cttttgtggt taaaagagac ctttgggtca gtctgcctca 60
ttccttgaag agtttagccc tggctcactt ttcactctat ttcttctcct gtctcaagaa 120
agaagaaaaa aagagacaaa ttaccagaa accctcctt tccccacatg gaggccttgg 180
caaatgttaa ttttcctaga aaatccttca gacctgaaga cgcaggaaaa gaatctggct 240
ctcagggtgg cttctgcgtc ccgcggcca ggccccagac tatggtcaca gggccgtcct 300
gttctctccc gggactccag aatttctctc ctcaaaggaa agaaaacagg gcatgcgctt 360
gttggcaaaa cgcaggggcg gctcccaaaa accccatgtg tgtacgatta aaagttggcc 420
gtccccaggc ctcccagcgc aaacttaaag agacagggct ttgctgaaaa ccaaactagg 480
gccagctggg ctttttaaca acctagagac tttccggagc tgcttgaac agagcctgcg 540
ggaaacgggg cttgccagag aactcacag tttccttcat ggctgtttt ggtcccctaa 600
gaatctccac atcattgtct ttcttgtgcc ttttcttgg tgagcaacag aaagggaagg 660
```

gttccaagcc tctaaaaatg tgctttgtga tcaggagtgc gctccaaacc aaatacgcgc 720
gctgcccttt cgaggccagt gagctcagcc tccaaggctt taaagccaca tttcagcaag 780
agaaagcgct gagagctcgc aggttcatta aagaaggcaa agcactgggt tctctcctta 840
gaaaagtagg tttcttggct tgatgtagac tggcttgctt tgatttttag tgaagggaat 900
gtacgtaaaa caaaataggg cttggctggt caaaggagac aagcaggatg gatggatgga 960
tggtatggatg gatgtatgga tgaatagata gatggtgtt gcatgtaaat tgcagagaaa 1020
acaaaaccaa agctgattgg aaacaattaa ttgtgggtgt ctgaggggga aggtcgcagc 1080
tttgggcagc tttgagaagc ggtacaagag ttctgtgcct gtgtgtccag ccctggagcc 1140
agccagtgca tttattttta gctcttagaa gcaactcctt ggcccaggaa tgcgtgaccc 1200
ctgagatggg tccacgcata tctctacact tcttctctc cgtgggatac tggactcgtg 1260
cctctgcgcc cattctcttc tcacgcata ccatgagctt taatttact ttctgatcac 1320
ggtacgtcga taaagccagt attacactta aatgaagtat tctttttgt aatcgttttt 1380
tttagaaggt aaacaaattt aataaagcta ccaataatga gaaaaaaaaa aaaaaaaaaa 1439

<210> 7

<211> 3047

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No.: 1656953CB1

<400> 7

cgagacagag gaaatgtgtc tccctccaag gccccaaagc ctcagagaaa ggggtgtttct 60
gggtttgcct tagcaatgca tcggtctctg aggtgacact ctggagcggg tgaaggcca 120
caaggtgcag ggttaatact cttgccagtt ttgaaatata gatgctatgg ttcagattgt 180
ttttaataga aaactaaagg ggcaggggaa gtgaaaggaa agatggaggt tttgtgcggc 240
tcgatggggc atttggaact tctttttaaa gtcactctcat ggtctccagt tttcagttgg 300
aactctgggt tttaacactt aaggagagac aaggctgtgt ccatttggca aaacttcctt 360
ggccacgaga ctctaggtga tgtgtgaagc tgggcagctc gtggtgtgga gagcagccat 420
ctgtctggcc attcagagga ttctaaagac atggctggat gcgctgctga ccaacatcag 480
cacttaataa aatgcaaag caacatttct cctctgggc cttgaaaatc cttgccctta 540
tcatttgggg tgaaggagac atttctgtcc ttggcttccc acagcccaa cgcagtctgt 600
gtatgattcc tgggatccaa cgagccctcc tattttcaca gtgttctgat tgctctcaca 660
gccagggccc atcgtctgtt ctctgaatgc agccctgttc tcaacaacag ggaggtcatg 720
gaacccctct gtggaaccca caaggggaga aatgggtgat aaagaatcca gttcctcaaa 780
accttccctg gcaggtctgg tccctctcct gctgggtggg gctttctctt gcacaccact 840
cccaccacgg ggggagagcc agcaacccaa ccagacagct caggttgtgc atctgatgga 900
aaccactggg ctcaaacacg tgctttattc tcctgtttat ttttctgtt actttgaagc 960
atggaaattc ttgtttgggg gatcttgggg ctacagtagt gggtaaaca atgcccaccg 1020
gccaagaggc cattaacaaa tcgtccttgt cctgaggggc ccagcttgc tcgggcgtgg 1080
cacagtgggg aatccaaggg tcacagtatg gggagaggtg caccctgcc cctgctaact 1140
tctcgtaga cacagtgttt ctgccaggt gacctgttca gcagcagaac aagccagggc 1200
catggggacg ggggaagttt tcaactggag atggacacca agacaatgaa gatttgttgt 1260
ccaaataggt caataattct gggagactct tggaaaaaac tgaatatatt caggaccaac 1320
tctctccctc ccctcatccc acatctcaaa gcagacaatg taaagagaga acatctcaca 1380
caccagctc gccatgccta ctcatctctg aatttcaggt gccatcactg ctctttcttt 1440
cttctttgtc atttgagaaa ggatgcagga ggacaattcc cacagataat ctgaggaatg 1500
cagaaaaacc agggcaggac agttatcgac aatgcattag aacttgggtga gcatcctctg 1560
tagagggact ccaccctgc tcaacagctt ggcttccagg caagaccaac cacatctggg 1620
ctctgccttc ggtggccac acacctaagc gtcactgca ttgccatagc atcatgatgc 1680
aacacatcta cgtgtagcac tacgacgtta tgtttgggtg atgtggggat gaactgcatg 1740
aggctctgat taaggatgtg gggaagtggg ctgcggtcac tgtcggcctt gcaaggccac 1800
ctggaggcct gtctgttagc cagtgggtga ggagcaaggc ttcaggaagg gccagccaca 1860
tgccatcttc cctgcgatca ggcaaaaaag tgaattaaa aagtcaaacc tttatatgca 1920
tgtgttatgt ccattttgca ggatgaactg agtttaaaag aatttttttt tctcttcaag 1980
ttgctttgtc ttttccatcc tcatcacaag cccttgtttg agtgtcttat ccctgagcaa 2040
tctttcgtat gatggagatg atcattaggt acttttgttt caacctttat tcctgtaaat 2100
atcttctgtg aaactaggag aacagagatg agatttgaca aaaaaaaatt gaattaaaaa 2160
taacacagtc tttttaaaac taacatagga aagcctttcc tattatttct cttcttagct 2220
tctccattgt ctaaatacagg aaaacaggaa aacacagctt tctagcagct gcaaaatggg 2280

ttaatgcccc	ctacatattt	ccatcacctt	gaacaatagc	tttagcttgg	gaatctgaga	2340
tatgatccca	gaaaacatct	gtctctactt	cggctgcaaa	acccatgggt	taaactctata	2400
tggtttgtgc	atcttctcaa	ctaaaaatag	agatgataat	ccgaattctc	catatattca	2460
ctaatacaag	acactatctt	catactagat	tcctgagaca	aatactcact	gaagggcttg	2520
tttaaaaata	aattgtgttt	tggctctgtt	ttgtagataa	tgcccttcta	ttttaggtag	2580
aagctctgga	atccctttat	tgtgctgttg	ctcttatctg	caaggtggca	agcagttctt	2640
ttcagcagat	tttgccctact	attcctctga	gctgaagttc	tttgcataga	tttggcttaa	2700
gcttgaatta	gatccctgca	aaggcttgct	ctgtgatgtc	agatgtaatt	gtaaatgtca	2760
gtaatcactt	catgaacgct	aaatgagaat	gtaagtattt	ttaaatgtgt	gtatttcaaa	2820
tttgtttgac	taattctgga	attacaagat	ttctatgcag	gatttacctt	catcctgtgc	2880
atgtttccca	aactgtgagg	aggggaaggct	cagagatcga	gcttctcctc	tgagttctaa	2940
caaaatgggt	ctgtgagggg	cagccttttag	gaaggtgcag	ctttgttgtc	ctttgagctt	3000
tctgttatgt	gcctatccta	ataaactctt	aaacacaaaa	aaaaaaa		3047

<210> 8

<211> 3017

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No.: 1662318CB1

<400> 8

cgcaaaactca	acccttttcgg	aaacaccttc	ctcaacaggt	tcatgtgtgc	ccagctccct	60
aatcaggtcc	tggagagcat	cagcatcatc	gacaccccg	gtatcctgtc	gggtgccaa	120
cagagagtga	gccgcggcta	cgacttcccc	gccgtgctgc	gctgggttcg	ggagcgcgtg	180
gacctcatca	tcctgctctt	tgatgcgcac	aagctggaga	tctcggacga	gttctcagag	240
gccatcgccg	cggtgcgggg	ccatgaggac	aagatccggg	tgggtgctca	caagggccgac	300
atggtggaga	cgcagcagct	gatgcgcgtc	tacggcgcgc	tcatgtgggc	gctgggcaag	360
gtggtgggca	cgcccgaggt	gctgcgcgtc	tacatcggtt	ccttctgggt	ccagcccctc	420
ctggtgcccg	acaaccggcg	cctcttcgag	ctggaggagc	aggacctctt	ccgcgacatc	480
cagggcctgc	cccggcacgc	agccttgccg	aagctcaacg	acctgggtga	gagggcccg	540
ctggtgcgag	ttcacgctta	catcatcagc	tacctgaaga	aggagatgcc	ctctgtgttt	600
gggaaggaga	acaagaagaa	gcagctgatc	ctcaaactgc	ccgtcatctt	tgcaagatt	660
cagctggaac	atcacatctc	ccctggggac	tttcttgatt	gccagaaaat	gcaggagctg	720
ctgatggcgc	acgacttcac	caagtttcac	tcgctgaagc	cgaagctgct	ggaggcactg	780
gacgagatgc	tgacgcacga	catcgccaag	ctcatgcccc	tgctgcggca	ggaggagctg	840
gagagcaccg	aggtgggcgt	gcaggggggc	gcttttgagg	gcacccacat	ggggccgttt	900
gtggagcggg	gacctgacga	ggccatggag	gacggcgagg	agggctcgga	cgacgaggcc	960
gagtgggtgg	tgaccaagga	caagtccaaa	tacgacgaga	tcttctacaa	cctggcgcct	1020
gccgacggca	agctgagcgg	ctccaaggcc	aagacctgga	tgggtggggac	caagctcccc	1080
aactcagtgc	tggggcgcat	ctggaagctc	agcgatgtgg	accgcgacgg	catgctggat	1140
gatgaagagt	tcgcgctggc	cagccacctc	atcgaggcca	agctggaagg	ccacgggctg	1200
cccgccaaac	tgccccgctg	cctggtgcca	ccctccaagc	gacgccacaa	gggctccgcc	1260
gagtgagccg	ggccccctc	ccatggccct	gctgtggctc	cccagctcca	gtcggctgca	1320
cgcacacccc	tgctccggct	cacacacgcc	ctgcctgccc	tccttgccca	gctgtaagga	1380
ccgggggtct	ccctcctcac	taccgccaga	caccccggtg	gaagcattta	gaggggacca	1440
cgggagggac	aaggcttctc	tgtccgccct	tcacacctcc	agcctcacgt	tcacttaggc	1500
acatcacaca	cacactggca	cacgcaggca	tccatccatc	cgtcattcat	tcaaataatt	1560
attgagcacc	tactatgtgc	ccagccctgt	tctaggcact	gggcattacc	atagagaaca	1620
aaatagacaa	atacatctgc	cctcatggaa	ggtgacgttc	ccaggagagg	gcacctacac	1680
agtcacgcaa	acacacacta	attcctggca	gggccccag	cccctcccct	ggctgagcag	1740
ccctgtggct	gaaatgacta	gcagataaac	agacccctt	ctgctccgct	tcctcctgcc	1800
cagccaggca	acaccctcaa	cggctccat	cacatcctca	ggtctcggga	ccatgggggg	1860
ctcagagggg	agacacacct	actgcttcc	cagatggggc	cctccgcagc	cccttccctt	1920
gctcggggaa	agcccccaat	tctgcccaca	cccatttatt	tccttccttc	cttcccttct	1980
ttctttcctt	ccttcccttct	ttttgtttt	tgcccccaat	tctgcccata	cccatttctt	2040
tctttccttc	cttcccttct	ttttgtttt	gccccagtt	ctgtccacac	cccttccctt	2100
tcctgtcctg	tcctttcttt	cttttttgat	agaatcttgc	tctgtcggcc	aggctgggag	2160
tgagtggtg	agatctcagc	tcactgcaac	ctccacctcc	tgggttgaag	tgattctcgt	2220
gcctcagcct	cctgagtagc	tgggactgca	ggcacgcgcc	accacgccc	gctaattttt	2280

```

gtatttgagt agagacgggg tttcaccatg ttggccaggc tggctctgaa ctccgcacat 2340
caggtgatct gctcgccctcg gcctcccaaa gtgatgggat tacaggcatg agccaccgtg 2400
cccggcttca caccatttc tttaaaaagg atcccgtagc aggcagaaaa gccccttcca 2460
tctgtctcct ctgatactgt gcccccttgg agatatttcc gtcctccacc cacgtgtctg 2520
tggctggaac tgcccagcct gctcctggcc ccctggaagc ctccccacag ctggtaatct 2580
ggacttaagg attgctgggc caccgcctct ctgcctacca ccattccata ttttaagtga 2640
gcccctacgt agaaaggccc cggggcttta ttttagtctc cttttcaggg atgtcgtggg 2700
cgggggaggg ggttcttggg gctacagccc tctccccacc cctaaaggga cgccgacgct 2760
gtttgctgcc ttcaccacat attagtgtt gaccctggca ggggaccca tggaaaagat 2820
ggggaagagc aaaatacatg gagacgacgc accctccagg atgtcgtctg ggattccac 2880
gcccaccact gtccccacc ccatggctgg gaggggctc tgaacggaac agtgtccca 2940
cagagcgaat aaagcaaggc ttcttcccca aaaaaaaaaa aaaaaaaaaa attggtgcgg 3000
ccgaagttat tcccttc                                     3017

```

<210> 9
 <211> 1735
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No.: 1996726CB1

```

<400> 9
tcgggaggaa ggagactaca cctgctttgc tgaaaatcag gtcgggaagg acgagatgag 60
agtcagagtc aagggtggtga cagcgcccg caccatccgg aacaagactt acttggcggt 120
tcaggtgccc tatggagacg tggctactgt agcctgtgag gccaaaggag aacctatgcc 180
caaggtgact tggttgtccc caaccaacaa ggtgatcccc acctcctctg agaagtatca 240
gatataccaa gatggcactc tccttattca gaaagccag cgttctgaca gcggaacta 300
cacctgcttg gtcaggaaca gcgcgggaga ggataggaag acggtgtgga ttcacgtcaa 360
cgtccagcca cccaagatca acggtaaccc caaccccatc accaccgtgc gggagatagc 420
agccgggggc agtcggaaac tgattgactg caaagctgaa ggcattccca ccccgagggt 480
gttatgggct tttcccagg gtgtggttct gccagctcca tactatggaa accggatcac 540
tgtccatggc aacggttccc tggacatcag gagtttgagg aagagcgact ccgtccagct 600
ggtatgcatg gcacgcaacg agggagggga ggccagggtg atcgtgcagc tcaactgtcct 660
ggagcccatg gagaaaccca tcttcacga cccgatcagc gagaagatca cggccatggc 720
gggccacacc atcagcctca actgctctgc cgcggggacc ccgacacca gcctggtgtg 780
ggtccttccc aatggcaccg atctgcagag tggacagcag ctgcagcgct tctaccacaa 840
ggctgacggc atgctacaca ttagcgggtc ctccctcggtg gacgccgggg cctaccgctg 900
cgtggcccg c aatgccgtg gccacacgga gaggtggtc tccctgaagg tgggactgaa 960
gccagaagca aacaagcagt atcataacct ggtcagcatc atcaatggtg agaccctgaa 1020
gctcccctgc acccctcccg gggctgggca gggacgttct tcttgacgc tccccaatgg 1080
catgcatctg gagggcccc aaacctggg acgcgtttct cttctggaca atggcaccct 1140
cacggttcgt gaggcctcgg tgtttgacag ggttacctat gtatgcagga tggagacgga 1200
atacggccct tcggtcacca gcatccccgt gattgtgatc gcctatcctc cccggatcac 1260
cagcgagccc acccgggtca tctacaccgg gcccggaac accgtgaaac tgaactgcat 1320
ggctatgggg attcccaaag ctgacatcac gtgggagtta ccggataagt cgcattctgaa 1380
ggcaggggtt caggctcgtc tgtatggaaa cagatttctt cacccccagg gatcactgac 1440
catccagcat gccacacaga gagatgccgg cttctacaag tgcattggaa aaaacattct 1500
cggcagtgac tccaaaacaa cttacatcca cgtcttctga aatgtggatt ccagaatgat 1560
tgcttaggaa ctgacaacaa agcgggggtt gtaagggaag ccaggttggg gaataggagc 1620
tcttaataaa tgtgtcacag tgcattggtg cctctggtgg gtttcaagtt gaggttgatc 1680
ttgatctaca attgttgga aaaggaagca atgcagacac gagaaggagg gctca 1735

```

<210> 10
 <211> 1016
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature

<223> Incyte ID No.: 2137155CB1

<400> 10

ctgtacgttc	ccctgtggcc	cacgcctagt	gaaaafgata	tcgtacatct	ccctagagat	60
atgggtcacc	tccaggtaga	ttacagagat	aacaggctgc	acccaagtga	agattcttca	120
ctggactcca	ttgcctcagt	tgtggttccc	ataattatat	gcctctctat	tataatagca	180
ttcctattca	tcaatcagaa	gaaacagtgg	ataccactgc	tttgctggta	tcgaacacca	240
actaagcctt	cttccttaaa	taatcagcta	gtatctgtgg	actgcaagaa	aggaaccaga	300
gtccaggtgg	acagttccca	gagaatgcta	agaattgcag	aaccagatgc	aagattcagt	360
ggcttctaca	gcatgcaaaa	acagaaccat	ctacaggcag	acaatttcta	ccaaacagtg	420
tgaagaaagg	caactaggat	gaggtttcaa	aagacggaag	acgactaaat	ctgctctaaa	480
aagtaaaacta	gaatttgtgc	acttgcttag	tggattgtat	tggattgtga	cttgatgtac	540
agcgctaaga	ccttactggg	atgggctctg	tctacagcaa	tgtgcagaac	aagcattccc	600
acttttcttc	aagataactg	accaagtgtt	tcttagaacc	aaagttttta	aagttgctaa	660
gatataatttg	cctgtaagat	agctgtagag	atatttgggg	tggggacagt	gagtttgat	720
ggcgaaatac	accgcacggg	ggtgttgagg	agaaaaattt	gtcagcttgg	ctcggggaga	780
aacctgtgta	cactaaagca	gttcagtgtg	ccagagggtta	tttttttccc	attgctctga	840
agactgcact	ggttgctgca	aactcaggcc	tgaatgagcg	gaaacaaaaa	aagccttgcg	900
ccccgatgcc	ataacacctt	tggaatcccg	agcggccctc	agaaaccttt	tcaggcatcc	960
aggtcttaag	cccaagtatc	tttctataca	gtcccactgc	ggtgagcgtg	ggggag	1016

<210> 11

<211> 2288

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No.: 2268890CB1

<400> 11

caaccagggt	caggctgtgc	tcacagtttc	ctctggcggc	atgtaaaggc	tccacaaagg	60
agttgggagt	tcaaagtagg	ctgctgcgga	cggcctgagg	atggacccca	agccctggac	120
ctgccgagcg	tggcactgag	gcagcggtcg	acgctactgt	gagggaaaga	aggttgtgag	180
cagccccgca	ggacccttgg	ccagccctgg	ccccagcctc	tgccggagcc	ctctgtggag	240
gcagagccag	tggagcccag	tgaggcaggg	ctgcttgcca	gccaccggcc	tgcaactcag	300
gaaccctctc	agaggccatg	gacaggctgc	cccgctgacg	gccagggtga	agcatgtgag	360
gagccgcccc	ggagccaagc	aggagggaag	aggctttcat	agattctatt	cacaaagaat	420
aaccaccatt	ttgcaaggac	catgaggcca	ctgtgcgtga	catgctgggt	gctcggactg	480
ctggctgcca	tgggagctgt	tgcaggccag	gaggacggtt	ttgagggcac	tgaggagggc	540
tcgccaagag	agttcattta	cctaaacagg	tacaagcggg	cgggcgagtc	ccaggacaag	600
tgcacctaca	ccttcattgt	gccccagcag	cgggtcacgg	gtgccatctg	cgtcaactcc	660
aaggagcctg	aggtgcttct	ggagaaccga	gtgcataagc	aggagctaga	gctgctcaac	720
aatgagctgc	tcaagcagaa	gcggcagatc	gagacgctgc	agcagctggt	ggagggtggac	780
ggcggcattg	tgagcgaggt	gaagctgctg	cgcaaggaga	gccgcaacat	gaactcgcgg	840
gtcacgcagc	tctacatgca	gctcctgcac	gagatcatcc	gcaagcggga	caacgcgttg	900
gagctctccc	agctggagaa	caggatcctg	aaccagacag	ccgacatgct	gcagctggcc	960
agcaagtaca	aggacctgga	gcacaagtac	cagcacctgg	ccacactggc	ccacaaccaa	1020
tcagagatca	tcgcgcagct	tgaggagcac	tgccagaggg	tgccctcggc	caggcccgtc	1080
ccccagccac	cccccgctgc	cccgccccgg	gtctaccaac	caccaccta	caaccgcac	1140
atcaaccaga	tctctaccaa	cgagatccag	agtgaccaga	acctgaaggt	gctgccaccc	1200
cctctgcccc	ctatgcccac	tctcaccagc	ctcccattct	ccaccgacaa	gccgtcgggc	1260
ccatggagag	actgcctgca	ggccctggag	gatggccacg	acaccagctc	catctacctg	1320
gtgaagccgg	agaacaccaa	ccgcctcatg	cagggtgtgt	gcgaccagag	acacgacccc	1380
gggggctgga	ccgtcatcca	gagacgcctg	gatggctctg	tttaacttct	caggaactgg	1440
gagacgtaca	agcaagggtt	tgggaacatt	gatggcgaat	actggctggg	cctggagaa	1500
atttactggc	tgacgaacca	aggcaactac	aaactcctgg	tgacctgga	ggactggtcc	1560
ggccgcaaag	tctttgcaga	atacgccagt	ttccgcctgg	aacctgagag	cgagtattat	1620
aagctgcggc	tggggcgcta	ccatggcaat	gcgggtgact	cctttacatg	gcacaacggc	1680
aagcagttca	ccaccctgga	cagagatcat	gatgtctaca	caaggaaactg	tgcccactac	1740
cagaagggag	gctgggtgga	taacgcctgt	gcccactcca	acctcaacgg	ggtctggtac	1800
cgcggggggc	attaccggag	ccgctaccag	gacggagtct	actgggctga	gttccgagga	1860

ggctcttact cactcaagaa agtgggtgatg atgatccgac cgaaccccaa caccttccac 1920
taagccagct cccctcctg acctctcgtg gccattgcca ggagcccacc ctgggtcacgc 1980
tgccacagc acaaagaaca actcctcacc agttcatcct gaggctggga ggaccgggat 2040
gctggattct gttttccgaa gtcactgcag cggatgatgg aactgaatcg atacggtgtt 2100
ttctgtccct cctactttcc ttcacaccag acagcccctc atgtctccag gacaggacag 2160
gactacagac aactctttct ttaaataaat taagtctcta caataaaaac acaactgcaa 2220
agtaccttca taatatacat gtgtatgagc ctcccttggtg cacgtatgtg tatagcacat 2280
atatatgg 2288

<210> 12

<211> 3304

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No.: 2305981CB1

<400> 12

ccctcttatg gattcccagc aagcatcagg aaccattgtg caaattgtca tcaataacaa 60
acacaagcat ggacaagtgt gtgtttccaa tggaaagacc tattctcatg gcgagtcctg 120
gcacccaaac ctccgggcat ttggcattgt ggagtgtgtg ctatgtactt gtaatgtcac 180
caagcaagag tgtaagaaaa tccactgccc caatcgatac ccctgcaagt atcctcaaaa 240
aatagacgga aagtgtgtgca aggtgtgtcc aggtaaaaaa gcaaaagaag aacttccagg 300
ccaaagcttt gacaataaag gctacttctg cggggaagaa acgatgcctg tgtatgagtc 360
tgtattcatg gaggatgggg agacaaccag aaaaatagca ctggagactg agagaccacc 420
tcaggtagag gtccacgttt ggactattcg aaagggcatt ctccagcact tccatattga 480
gaagatctcc aagaggatgt ttgaggagct tcctcacttc aagctgggtg ccagaacaac 540
cctgagccag tggaaagtct tcaccgaagg agaagctcag atcagccaga tgtgttcaag 600
tcgtgtatgc agaacagagc ttgaagattt agtcaagggt ttgtacctgg agagatctga 660
aaagggccac tgtaggcaa gacagacagt attggatagg gtaaagcaag aaaactcaag 720
ctgcagctgg actgcaggct tattttgctt aagtcaacag tgccctaaaa ctccaaactc 780
aatgcagtc aattattcac gccatgcaca gcataatttg ctcttttgtg tggagtgggtg 840
tgtcagccct tgaacatctc ctccaaagag actagaagag tcttaaatta tatgtgggag 900
gaggagggat agaacatcac aacactgctc tagtttcttg gagaatcaca tttctttaca 960
ggttaaagac aaacaagacc ccagggtttt tatctagaaa gttattcaag tgaaagaaag 1020
agaagggaat tgcttagtag gagttctgca gtatagaaca attacttgta tgaaattata 1080
cctttgaatt ttagaatgtc atgtgttctt ttaaaaaaat tagctcccca tcctccctcc 1140
tactccctc cctccctcct tctctctctc tctctctctc cctctctcac agacacacac 1200
acacacacac acacacgcac acgcacgtcc acactcacat taaactaaag ctttatttga 1260
agcaaagcta gccaaaattc tacgttactt ttcccttgac tggatcccaa gtagcttgga 1320
agtttttgtg ccaggagag taaataactg tgaacaagag gctctgccct taggtctttg 1380
tggtgtttta agtcaccaac aatagagtca gggtaaagaa taaaaacact ttcatagcct 1440
cattcattca cttagaagtg gtaataattt ttccctaatt ataccacttt tcttttcccc 1500
ctgtacctat gggacttcca gaaagaagtt aaattgagta aaatcatcag aaactgaatc 1560
catgtaagaa aaaataattg ttgaagaaag aagttgatag aattcaaaaa ggccatcttt 1620
ttgctttcac atcaataaaa tttaccaagt aatagatcag tactactaa tatttttgag 1680
accatagttg tctggtcaga aaaattatat taaattagta aattctagaa gctctttaa 1740
aggggaagttt tccttcttct ccaattatag gagttgattt ttactttgca aagtggctcg 1800
gtcctcatga gcatctgcat gttgactctt cagtttaagaa aattgttggt catttaggga 1860
ggtggatatt ctgatgaaga tctttatcct aaaccttcct actatccttg tcttattcat 1920
caagcagata ttttagtcaa gaattccaga gaaggctgct cctaaaaatgt ctacttgag 1980
cccaatacca gagcataaac tatccattct ggggtctggc tttagaaatc atctttgtgg 2040
gaagacctaa ttcttcacag caaggatctc aggcattgct tctagatttg ttccctctga 2100
ggggcaggaa tgaactgtag aaatgtttta aggaccaga aaccctatat gtctcattcc 2160
atgactatag gtgagagaat tctttcctaa gaggtttga taccaatagg ggaaaatgta 2220
aaatgttcag tctttatgac aacctggcat aaaggagtca attcttatga aagagacaca 2280
agggccttat ggccagggtt tcttgggaca agactctcac cagcacatca cacacgttct 2340
ccttgaaga gagaagcagt acatcccggg tgagaggtca caaagcatta gtttgtgtgt 2400
gtgtgtgtgt gtgtgtgtgt gtgtgtgtgt gtgtgtgtgt gtggtaaagg ggggaagggtg 2460
ttatgcggct gctccctccg tcccagaggt ggcagtgatt ccataatgtg gagactagta 2520
actagatcct aaggcaaaga ggtgtttctc cttctggatg attcatccca aagccttccc 2580

```

accaggtgt tctctgaaag cttagcctta agagaaacacg cagagagttt ccctagatat 2640
actcctgcct ccaggtgctg ggacacacct ttgcaaaatg ctgtgggaag caggagctgg 2700
ggagctgtgt taagtcaaag tagaaaccct ccagtgtttg gtgttggtga gagaatagga 2760
catagggtaa agaggccaag ctgcctgtag ttagtagaga agaattggatg tggttcttct 2820
tgtgtattta tttgtatcat aaacacttgg aacaacaaag accataagca tcatttagca 2880
gtttagacca ttttctagtt aactcatgta aacaagtaag agtaacataa cagtattacc 2940
ctttcactgt tctcacagga catgtaccta attatggtag ttatttatgt agtcactgta 3000
tttctggatt tttaaattaa taaaaaagtt aattttgaaa aaaaaaaaaa aaaaaaaaaa 3060
aaaaaaaaaa aaaaaaaaaa actcgagggg gggcctgtac cgggttcccc gtaacaggtt 3120
cgcccttaag attccctggc cgcagttttt ggccgcgttt tggggaacct ctgggtaccc 3180
ccttagttgc tcgctaaaat cccctttcgc agcccgttta aaggctgggg ccggccgatt 3240
gccttcccaa tagcctccca tgaatgggaa tgggaattgga agggaaattt tggtaaatcc 3300
ggta 3304

```

<210> 13
 <211> 708
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No.: 2457612CB1

```

<400> 13
ggaaagccag gaagtgcagg aatcatttca tcagggccaa taactacacc acccctgagg 60
tcaacaccca ggctactgg aactcccttg gagagaatag agacagatgt aaagcaacca 120
acagttcctg cctctggaga agaactggaa aatataactg actttagctc aagcccaaca 180
agagaaactg atcctcttgg gaagccaaga ttcaaaggac ctcatgtgcg atacatccaa 240
aagcctgaca acagtccctg ctccattact gactctgtca aacggttccc caaagaggag 300
gccacagagg ggaatgccac cagcccacca cagaaccac ccaccaacct cactgtggtc 360
accgtggaag ggtgcccctt catttgtcat cttggactgg gaaaagccac taaatgacac 420
tgtcactgaa tatgaagtta tatccagaga aaatgggtca ttcagtggga agaacaagtc 480
cattcaaatg acaaatcaga cattttccac agtagaaaat ctgaaaccaa acacgagtta 540
tgaattccag gtgaaaccca aaaaccgct tgggtgaaggc ccggtcagca acacagtggc 600
attcagtact gaatcagcgg acccagagtg agtgagcagt ttctgcagga gagatgcctc 660
tggaactgaag gccgctttgt tcgactcttg ctcaggtgta agggcaac 708

```

<210> 14
 <211> 2040
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No.: 2814981CB1

```

<400> 14
cggccagccg ccgcgcgctg cagctctccg ggacgcccgt gcgccagctg cagaagggcg 60
cctgcccgtt gggctctccac cagctgagca gccgcgcta caagttcaac ttcattgctg 120
acgtgggtga gaagatcgca ccagccgtgg tccacataga gctcttctg agacacccgc 180
tgtttggccg caacgtgccc ctgtccagcg gttctggctt catcatgtca gaggccggcc 240
tgatcatcac caatgcccac gtggtgtcca gcaacagtgc tgccccgggc aggcagcagc 300
tcaaggtgca gctacagaat ggggactcct atgaggccac catcaaagac atcgacaaga 360
agtcggacat tgccaccatc aagatccatc ccaagaaaaa gctccctgtg ttgttgctgg 420
gtcactcggc cgacctgcgg cctggggagt ttgtgggtggc catcggcagt cccttcgccc 480
tacagaacac agtgacaacg ggcacgtca gcaactgcca gcgggagggc agggagctgg 540
gcctccggga ctccgacatg gactacatcc agacggatgc catcatcaac tacgggaact 600
ccgggggacc actggtgaac ctggatggcg aggtcattgg catcaacacg ctcaaggtca 660
cggctggcat ctctttgcc atcccctcag accgcacac acggttcctc acagagttcc 720
aagacaagca gatcaaagac tggagaagc gcttcacatg catacgatg cggacgatca 780

```

caccaagcct ggtggatgag ctgaaggcca gcaacccgga cttcccagag gtcagcagtg 840
gaatttatgt gcaagaggtt gcgccgaatt caccttctca gagaggcggc atccaagatg 900
gtgacatcat cgtcaagggtc aacgggcgtc ctctagtga ctcgagtga ctgcaggagg 960
ccgtgctgac cgagtctect ctectactgg aggtgcggcg ggggaacgac gacctectct 1020
tcagcatcgc acctgaggtg gtcattgtgag gggcgcatte ctccagcgcc aagcgtcaga 1080
gcctgcagac aacggagggc agcgccccc cgagatcagg acgaaggacc accgtcggtc 1140
ctcagcaggg cggcagcctc ctctgggtg tccggggcag agcgagggtt gggcttggcc 1200
aggggcccga atttccgctt ggggagtggt ggatccacat cccggtgccg gggagggaag 1260
cccaacatcc ccttgtacag atgatcctga aagtcacttc caagttctcc ggatattcac 1320
aaaactgcct tccatggagg tcccctctc tctagtctc ccgctctgc cctgtgaac 1380
acctatctgc agtatccctt gctcctgccc ctctactgc aggtctgggc tgccaagctt 1440
cttccccctt gacaaacgcc cacctgacct gaggccccag cttccctctg ccttaggact 1500
taccaagctg tagggccagg gctgctgcct gccagcctgg ggtccctgga ggacaggtca 1560
catctgatcc ctttggggtg cgggggtggg gtccagccca gagcaggcac tgagtgaatg 1620
ccccctggct gcggagctga gcccgcctt gccatgaggt tttcctcccc aggcaggcag 1680
gaggccgagg ggagcacgtg gaaagtggc tgctgcctgg ggaagcttct cctccccaag 1740
gcggccatgg ggcagcctgc agaggacagt ggacgtggag ctgcggggtg tgaggactga 1800
gccggcttcc ccttcccacg cagctctggg atgcagcagc cgctcgcatg gaagtgccgc 1860
ccagaggcat gcaggctgct gggcaccacc cctcatcca gggaacgagt gtgtctcaag 1920
gggcatttgt gagctttgct gtaaatggat tcccagtggt gcttgtactg tatgtttctc 1980
tactgtatgg aaaataaagt ttacaagcac aaaaaaaaaa aaaaaaaaaa aaaaaaaagg 2040

<210> 15

<211> 2121

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No.: 3089150CB1

<400> 15

gtaaaagctg gttgtgatcg catcatagac tccaaaaaga agtttgataa atgtggtggt 60
tgccgggggaa atggatctac ttgtaaaaaa atatcaggat cagttactag tgcaaacct 120
ggatatcatg atatcatcac aattccaact ggagccacca acatcgaagt gaaacagcgg 180
aaccagaggg gatccaggaa caatggcagc tttcttgcca tcaaagctgc tgatggcaca 240
tatattctta atggtgacta cactttgtcc accttagagc aagacattat gtacaaagg 300
gttgtcttga ggtacagcgg ctctctgctg gcattggaag gaattcgag ctttagccct 360
ctcaaagagc ccttgaccat ccaggttctt actgtgggca atgcccttcg acctaaaatt 420
aaatacacct acttcgtaaa gaagaagaag gaatctttca atgctatccc cactttttca 480
gcatgggtca ttgaagagtg gggcgaatgt tctaagtcag gtgaattggg ttggcagaga 540
agactggtag aatgccgaga cattaatgga cagcctgctt ccgagtgtgc aaaggaagt 600
aagccagcca gcaccagacc ttgtgcagac catccctgcc cccagtggca gctgggggag 660
tggtcatcat gttctaagac ctgtgggaag ggttacaata aaagaagctt gaagtgtctg 720
tcccatgatg gaggggtggt atctcatgag agctgtgatc ctttaaagaa acctaaacat 780
ttcatagact tttgcacaat ggcagaatgc agttaagtgg ttttaagtgg gttagctttg 840
agggcaaggc aaagtgagga agggctggtg cagggaaagc aagaaggctg gagggatcca 900
gcgtatcttg ccagtaacca gtgaggtgta tcagtaagg gggattatgg gggtagatag 960
aaaaggagtt gaatcatcag agtaaaactgc cagttgcaaa tttgatagga tagttagtga 1020
ggattattaa cctctgagca gtgatatagc ataataaagc cccgggcatt attattatta 1080
tttcttttgt tacatctatt acaagtttag aaaaaacaaa gcaattgtca aaaaaagtta 1140
gaactattac aaccctgtgt tcctgggtact tatcaaatac ttagtatcat gggggttggg 1200
aaatgaaaag taggagaaaa gtgagatttt actaagacct gttttacttt acctactaa 1260
caatgggggg agaaaggagt acaaatagga tctttgacca gcactgttta tggctgctat 1320
ggtttcagag aatgtttata cattatttct accgagaatt aaaacttcag attgttcaac 1380
atgagagaaa ggctcagcaa cgtgaaataa cgcaaatggc ttctctcttc cttttttgga 1440
ccatctcagt ctttatttgt gtaattcatt ttgaggaaaa aacaactcca tgtatttatt 1500
caagtgcatt aaagtctaca atggaaaaaa agcagtgaag cattagatgc tggtaaaagc 1560
tagaggagac acaatgagct tagtacctcc aacttccttt ctttctacc atgtaaccct 1620
gctttgggaa tatggatgta aagaagtaac ttgtgtctca tgaaaatcag tacaatcaca 1680
caaggaggat gaaacgccgg aacaaaaatg aggtgtgtag aacagggtcc cacaggtttg 1740
gggacattga gatcacttgt cttgtggtgg ggaggctgct gaggggtagc aggtccatct 1800

ccagcagctg gtccaacagt cgtatcctgg tgaatgtctg ttcagctctt ctgtgagaat 1860
atgatttttt ccatatgtat atagtaaaat atgttactat aaattacatg tactttataa 1920
gtattgggtt ggggtgttcct tccaagaagg actatagtta gtaataaatg cctataataa 1980
catattttatt tttatacatt tattttctaata gaaaaaaact tttaaattat atcgcttttg 2040
tggaagtgcata taaaaaatag agtattttata caatatatgt tactagaaat aaaagaacac 2100
ttttggaaaa aaaaaaaaaa a 2121

<210> 16

<211> 2900

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<223> Incyte ID No.: 3206667CB1

<400> 16

gaagtttttaa aaaaaactac agcagccaaa gaaactatat atatatatat atatatccag 60
aatgattgcc tctactgtcc tcattgactt gtttgaacct tagtgcctta ccctgtcctc 120
ttcccagttc tctttataga agctctagga gctttcgaaa agccaaagtc tttctgaaga 180
atctgtgctg gacagacata attccctttc tcattgtctc catctttgtt ggtcatggta 240
aggtttttcc atcagcctct gaaaaaatag ttgtgcacaa catctgctca ctggactgtc 300
tgatccaatg taattggctg cgtctggcta attctaagca ctaaagtcta catctaagct 360
atagatttaa gcttgaagct acagattata tcaactatcac caccaccct caccagtgga 420
aatcagacag tcagtcactt taagttaaag atatttgttg tctttgaatg atttgctgtc 480
acagactatt tggtagaaga aatatttttc acctgagaga ggaagagaaa tttctctagt 540
aacacaaaga gtgagttcta aaaggcatgc ccacatctct ttcgtgcctt aaggatagtg 600
agatgcacac ttatatatat actgtatata tttatatatt tatatatata tttcatatat 660
atatataata ttgcaagctt aagtttgcaa tttcccaaac aatacaaaaa gcaaattaca 720
caccctcacc actgttctta tctctatagt gatgaaacat taattaggga tcttgctgct 780
tttctttttc tacacgaagt tttcattaaa gccacagaat aattgatagg gcagctgttt 840
gagaacaggt cccattttca cattagggct ttaaataaat tagaaactat ttgaggctat 900
aaaaatgtcc ttgagtttg agcctgagct ctgggtgaaat gctgatacat ctgatctatc 960
atgggaattg cagtttagaga gagtaaggaa taccatttag tcatctatcc gttcttcact 1020
tagcaggaat atgaaagaaa ggcacatgtt taagaggaat acctaaaggt ttttctaaat 1080
tccaacattt aaaaggcaat tgtgggctat ttttattttt taatattttg aaataaagtt 1140
tagtgtctag ggctgggagc caggactgat cttccatttc tttttctttg ttcccagcca 1200
tgcttttgta acttgccagg tggacttgac caactacatt acctgctgt gcctcagttt 1260
acctatttgt aaaatgggat taataatact tacctacctc acaggggtgt tgtgaggctc 1320
tattcatttg ctcttttatt ctttctgtga ttctctgtat gtccagcact ttgtagccat 1380
gggaggaaag ggactataaa agtgtacaat gttaatggaa tgatacggta cctgaaagcc 1440
ttgttttcta gtaagaaaat gctaccttgc tgtacatact tataaccttg tattttgaaa 1500
tgagaaatag gtttatattt tcagatctct caaaaatcac atcatttgac caaagaataa 1560
tttaagacac atagaacaga tttttttaat ttatatattt atcctgacca gcttagttct 1620
aataattttt agttgtgagt gattaaaaaa ctttggatca atttttgtca aacatgccaa 1680
ctttgtagtc tgagtgcag gcaaggattt ttgggtttta gatgcacttt tagcacacat 1740
ttgtatttcc cttggcatat cagattgagc taatgggtgat gttatttcaa tctaacagcc 1800
accaatctga aattgtattt caaatgttga ttctgtagtt ctttaaataa taatgaagct 1860
catcttatac attttgcttt caccaattga ttcttcttct ttttagccca ctattaaaac 1920
atttcttact gaatgggttca tgtaggcttg ctgaacagca cgcattactt gcttctgaa 1980
gagttcccc attcatccat ttgtcccat agttgctgtg gattatcaag ttttgaagga 2040
actgtacatc ccaacagact gaaacattct aagtgaatg agtataatcc aagtaactgg 2100
tgaacttttg aggtttggag cttgaagaga atggctaaga agatttgaat tatagggagg 2160
gaacagaaat catacatgaa aaggttttac tgagaagggg aaaaccttag atagagggac 2220
atgtgaaaca aatcatttg aaattttgat tcagacatcc atttccagt gcaaacagca 2280
aagcctgaac ccataaaccc aaatgatagg tgaagtggg tggttttatc caatgtctca 2340
agcaagcaat gtctgggaat atcatagagt aacaagtgt ggtcagccaa agaaacattc 2400
actgctggtg aaccaatacc ataagcatgt attatctaag cacttgatca agaaatatac 2460
atgttgatca agctctcaat tttgttcatt tattatcaaa tttttaaaat acaagtttgg 2520
tatgtgattt ggaaaagatg ctttctggat ctttaagccag ttgtcagtg aggtcctcag 2580
ggctgcaaat gtcaagacat aacctgttc ctcaccatca tgataccaga tacaggtgaa 2640
tacataggaa ctatctgcct gtgtcctcaa tctcccttca aacaagatgc tgattttag 2700

ggtacttggc aggttaaatt aaaccagaag aggtgactta ataaaaaagg gaatgacatt 2760
taggggtataa agatctcata agaaatgtaa tatgtaaatt atatcttgct ttatgttgta 2820
aaatatacat tgtttgcgct agaatagaaa tgatttcttt tcaataaaaa gaaagaagga 2880
ctctaaaaaa aaaaaaaaaa 2900

<210> 17

<211> 2507

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No.: 3284695CB1

<400> 17

cagagtgaag cttgtgcctg gtgaccaaag tccctccaaa gtgctcttcc ttctgggtta 60
ttcaagccaa atatctgggt ttccccctct cctcattccc tagcaaacc caattatctt 120
ccaagatagg agatatttcc catccccctc ctttgtaaatt atctcatctc ccactggaga 180
gccaggagc ctattcctgg catggatgtt ctgtccacac ttgaggctgg gcggtgtatc 240
agacccttca agcagcctgg ctggggccca ggactgagtc tggggtcagc ttccacggtc 300
gcttttccct tcctcaccac ccaccacagc ccacettgca tgcattggca gcccctccac 360
tccagcctga gccatgtgtg cccctgcggg aggaccatt catgccagaa agctggtaac 420
tccctcccag catccctgcg gaaggagtca gtttctgaga gtgtgacttt tcaaggcgaa 480
tgatggggaa ggggttccca gtccccacag tggccccacc tctggggcct gcaccagagc 540
ccttctgtgt cacggcgggc tgtgcaccca tgcacacacc tacgcacaca caacactccg 600
cactgcagta tattcttgcc aaagatttcc tttaaaagca agcactttta ctaattatta 660
ttttgtaaat gtttatcttc ttctgtcttc tccctccctg aatctatttt actgttgttt 720
attgttgaat ctgtgtgtca gccaggagag cgctgtctgg ccttgaacat gggctgggat 780
gggaaagggt ctgggagaag atgggcaaca aagagccagg gagtcatgga catcgagcgc 840
acgcagaccc cagcaggttc agtcccgtgc tggccaccagc tgtccagctg ggtgtctgga 900
gggaagaggg cagaggaggg tcatgtccct tcagctgggg gaggggcca gtgagctcca 960
cgtggctttt tcccaaaggg agcaagaggg aaggattggg cgagaaaaca atggagaggg 1020
gacctgcgaa ggaaaacagg gaggaagtga gcggtttgat cagcctgcta tcacgggtgtt 1080
ctggctctct tatttagcca ggcgcttaag ggacagatac atcacatcct aagtttgga 1140
aaggcctttg acccatgtca tctgagcgtc tcctccagta gctctgaaag ctgtggacac 1200
caatggccag gattccttct cccctggttt ttgaggatcc ctgggtcttc tgagactggc 1260
caggagaggg atgggtgggc cagtgggtgt gtgaaagcag gaggggcagc cctcctggac 1320
aagtgtgatc cccctataaa cggctctcag gaggttagtg agtaggagat tctgccttgt 1380
tctgatgagc ctgtgcaggg gctccagggg agcatgctgt ccagggggca cagaagggtg 1440
gtgagtgtga tcaaacttag tctcactccc acttttttag tctcactcct acttttgtcc 1500
accacccctg cctcctggat cttctccac tttttttt agcttttagga cctggggaga 1560
tcctgtgagt caaggcagac acccaatcct gccccacac tgggggtcct ccaagaggtt 1620
ggggggcaga gtcccagagc agccctttac cccagggtcca ggcctggaa tcctgagact 1680
cgcggtttct tggccagtgg taacacagga cgtgtgtgcg catgtgcaag tgtggatgta 1740
tgtgtgtgcg tgtgttttgc tcatttcttt agggaaactg ggagtcgggg ttggaggtgc 1800
tgggcaatgg aacttcaaat tcaatgtcgc ccagcagtga ggggagtcgg gaggtgaggc 1860
ctgtaggcca accaattggt ggagtctcag cgatagccca ggtgagaagt ggttcacca 1920
gaggggcagg gtgggggcct cgggcagatc tgtccctctt ggcccctctg tcctcaaatg 1980
tccaaaatgt tggaggacct ctgttcatat cccacgcctg ggctcttgcc agcagtggag 2040
ttactgtaga gggatgtccc aagcttggtt tccaatcagt gttaagctgt ttgaaactct 2100
cctgtgtctg tgttttggtt gtgcgtgtgt gtgagagcac atcagtgtgt gcaggctgtg 2160
tttccccatt tctctcctcc cttcagaccc atcattgaga acaaatgtaa gaaatccctt 2220
cccaccaccc tcctgcctc ccaggccctc tgcgggggaa acaagatcac ccagcatcct 2280
tccccacccc agctgtgtat ttatatagat ggaaatatat tttatatatt gtatcatcgt 2340
gcctatagcc gctgccaccg tgtataaatc ctgggtgtat ctccttatcc tggacatgaa 2400
tgtattgtac actgacgcgt cccactcct gtacagctgc tttgtttctt tgcaatgcat 2460
tgtatggctt tataaatgat aaagttaaag aaaaaaaaaa aaaaagg 2507

<210> 18

<211> 2929

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No.: 3481610CB1

<400> 18

```
aagctcggaa ttgggctcga gatgggttcc tcatcccttc ctgctgcaaa agaagttaac 60
aaaaaacaag tgtgctacaa acacaatttc aatgcaagct cagtttcctg gtgttcaaaa 120
actgttgatg tgtgttgta ctttaccat gctgctaata attcagtctg gagcccatct 180
atgaagctga atctgggtcc tggggaaaac atcacatgcc aggatcccgt aataggtgtc 240
ggagagccgg ggaaagtcac ccagaagcta tgccgggtct caaacgttcc cagcagccct 300
gagagtccca ttggcgggac catcacttac aaatgtgtag gctcccagtg ggaggagaag 360
agaaatgact gcatctctgc cccaataaac agtctgctcc agatggctaa ggctttgatc 420
aagagccctt ctcaggatga gatgtccctt acatacctga aggatctttc tattagcata 480
ggcaaagcgg aacatgaaat cagctcttct cctgggagtc tgggagccat tattaacatc 540
cttgatctgc tctcaacagt tccaacccaa gtcttgaaac cctggaaggt tttacaacag 600
tctacggtta atatcatcct tggcaagccc gtcttgaaac cctggaaggt tttacaacag 660
caatggacca atcagagttc acagctacta cattcagtg aaagattttc ccaagcatta 720
cagtcaggag atagccctcc attgtccttc tcccaaacta atgtgcagat gagcagcatg 780
gtaatcaagt ccagccaccc agaaacctat caacagaggt ttgttttccc atactttgac 840
ctctggggca atgtgggtcat tgacaagagc tacctagaaa acttgcagtc ggattcgtct 900
attgtcacca tggctttccc aactctccaa gccatccttg ctcaggatat ccaggaaaat 960
aactttgcag agagcttagt gatgacaacc actgtcagcc acaatacgac tatgccattc 1020
aggatttcaa tgacttttaa gaacaatagc ccttcaggcg gcgaaacgaa gtgtgtcttc 1080
tggaacttca ggcttgccaa caacacaggg ggggtgggaca gcagtgggtg ctatgttgaa 1140
gaaggtgatg gggacaatgt cactgtatc tgtgaccacc taacatcatt ctccatcctc 1200
atgtcccctg actccccaga tctagtttct ctctgggaa tactcctgga tattatttct 1260
tatgttgggg tgggcttttc catcttgagc ttggcagcct gtctagttgt ggaagctgtg 1320
gtgtggaaat cggtgaccaa gaatcggact tcttatatgc gccacacctg catagtgaat 1380
atcgctgcct cccttctggt cgccaacacc tgggttcattg tggctcgtgc catccaggac 1440
aatcgctaca tactctgcaa gacagcctgt gtggctgcca ccttcttcat ccacttcttc 1500
tacctcagcg tcttcttctg gatgctgaca ctgggcctca tgctgttcta tcgcctgggt 1560
ttcattctgc atgaaacaag caggtccact cagaaagcca ttgccttctg tcttggctat 1620
ggctgcccac ttgccatctc ggtcatcacg ctgggagcca cccagcccg ggaagtctat 1680
acgaggaaga atgtctgttg gctcaactgg gaggacacca aggcctgct ggctttcgcc 1740
atcccagcac tgatcattgt ggtggtgaac ataaccatca ctattgtgt catcaccaag 1800
atcctgaggg cttccattgg agacaagcca tgcaagcagg agaagagcag cctgtttcag 1860
atcagcaaga gcattggggc cctcacacca ctcttgggac tcaattgggg ttttgggtctc 1920
accactgtgt tcccagggac caacctgtg ttccatatca tatttgccat cctcaatgtc 1980
ttccagggat tattcatttt actctttgga tgcctctggg atctgaaggt acaggaagct 2040
ttgctgaata agttttcatt gtcgagatgg tcttcacagc actcaaagtc aacatccctg 2100
ggttcatcca cactgtgtt ttctatgagt tctccaatat caaggagatt taacaatttg 2160
tttggtaaaa caggaacgta taatgtttcc accccagaag caaccagctc atccctggaa 2220
aactcatcca gtgcttcttc gttgtcaaac taagaacagg ataatccaac ctacgtgacc 2280
tcccggggac agtggctgtg cttttaaaaa gagatgcttg caaagcaatg gggaacgtgt 2340
tctcggggca ggtttccggg agcagatgcc aaaaagactt tttcatagag aagaggcttt 2400
cttttgtaaa gacagaataa aaataattgt tatgtttctg tttgttccct cccctcccc 2460
cttgtgtgat accacatgtg tatagtattt aagtgaact caagccctca aggcccaact 2520
tctctgtcta tattgtaata tagaatttcg aagagacatt ttcacttttt acacattggg 2580
caciaagata agctttgatt aaagtagtaa gtaaaaggct acctaggaaa tacttcagtg 2640
aattctaaga aggaaggaag gaagaaagga aggaagaaag ggagggaac agggagaaag 2700
ggaaaaagaa gaaaaagaga tagatgataa taggaacaaa taaagacaaa caacattaag 2760
gggcatattg taagatttcc atgttaatga tctaataata tcaactcagtg ccacattttg 2820
agaatttttt tttttaatgg gcttcaaaaa ttggaaaact gtgaaagcta agtccattgg 2880
ggggaatgga attacttttg ggggccagta tctttccttt gattgttcc 2929
```

<210> 19

<211> 1725

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No.: 3722004CB1

<400> 19

```
gaggcaagaa ttcggcacga gggagagccc gcgggcgtgg gggagctcgg ggacctgcgg 60
accgggggag cccgaacgag ggggatcccc cgccggcgcc agcgaggcgg aggagcaggc 120
ggtggaggcg aggcaggaag aggagcagga cttggatggg gagaaggggc catcatcgga 180
agggcctgag gaggaggacg gagaaggctt ctccttcaaa tacagccccg ggaagctgag 240
gggaaaccag tacaagaaga tgatgaccaa agaggagctg gaggaggagc agaggattga 300
gctgacctct gacctcactt ccctgtagca agttccttag gtcctgagcc acaaatattc 360
ttgcaaatac ttttgaactg aagaataacg aagttatcct tagcgtcctc ctaaaggctt 420
ttccttttgg catcttaaaa gcttgagaga taaaacggaa accccagaga ggagtctggg 480
caggctccca gggtagcatg tgcctccata aatctgctga gctctagacc ctcaatcagg 540
acttgctcct tggctagcag gatcctggga acacctttgg ccctgccctg ttagagatg 600
ttcatgtctg ttctgtggg tcactttgtt aagctgaaga gttttaagag gtagagctca 660
gacctggac tgggattttt cttaccactc aaacttgcta tccacacacc ctgcacacct 720
tagataaaaa gaacatttta aaagcagagt tcactttcac tccagtctcc cctcttttgc 780
cctcactgaa gccaaaccac agaagacttt gaggaatgag agacaaatga ggtagagctc 840
acctgtgctc accagctccg tcagggtggg cagccgaccc ctttccttgg gaacccact 900
tctctctgtg gctggcttgg ttgtcggggg tgagatgcc aattgattac agggcagcaa 960
agaaccagta ccaggaattt acttgaccat tccccttatt ttcatctag aggaatctcg 1020
gattcagccc tttcattgct aagacacctt ttcactgagg ttcttaccag ctcagccaaa 1080
tctccactct gctatagcag aagcaataat gtttgcttta aaaagatttc ttgacctatg 1140
cctttttctt gaaagtittg tagattagtt agaacttcag atcatcagat cagtctcaaa 1200
tgggtttctt ggaattttat atttgacaat atttatacta taccaaactc atttgacgtt 1260
cttaggtttg ttggttaaaa cattttttta aagcagtaag tttatagaaa atgttttcat 1320
ttaatggaag gctggggaat gtccagcatc aaccctatg gcatgcattc ccagtggcct 1380
tctcatctgg gcctggaacc tttggttcag ggcttagggg agaacaggcc acatggcaac 1440
agccacacag tcattgcctt caacacagag ccacgtgtcc ccaaacagca atagtcatgc 1500
ccttgctccg gctgggatct aattgatata ataggtcggt gactccctcc tagtagagct 1560
atctaggttt gtctggaaag tttccgaccc tggcttatag gcaccacacc tcatgtactc 1620
ctcatggctt ggatctctgt attcagcctt tgttcagtc aataaacttt gagtagatga 1680
tctcaaaaaa aaaaaaaaaa aggccggcgc aagcttattc ctttt 1725
```

<210> 20

<211> 1987

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No.: 3948614CB1

<400> 20

```
gacggccagt gcaagctaaa attaaccctc actaaaggga ataagcttgc ggccgcctgg 60
agctctcgcc ctcggcttcg acgacggcaa cttctcgtcg ctcatccgcg cggtaggagga 120
gacggacgcg gggctgtaca cctgcaacct gcaccatcac tactgccacc tctacgagag 180
cctggccgtc cgcctggagg tcaccgacgg ccccccggcc accccgcct actgggacgg 240
cgagaaggag gtgctggcgg tggcgcgcg cgaccccgcg cttctgacct gcgtgaaccg 300
cgggcacgtg tggaccgacc ggcacgtgga ggaggtcaa caggtggtgc actgggaccg 360
gcagccgccc ggggtcccg acgaccgcgc ggaccgcctg ctggacctct acgcgtcggg 420
cgagcgccgc gcctacgggc ccttttttct gcgcgacgc gtggctgtgg gcgcggatgc 480
ctttgagcgc ggtgacttct cactgcgtat cgagccgctg gaggtcgccg acgagggcac 540
ctactcctgc cacctgcacc accattactg tggcctgcac gaacgcgcg tcttccacct 600
gacggtcgcc gaacccacg cggagccgcc cccccgggc tctccgggca acggctccag 660
ccacagcggc gccccaggcc cagacccac actggcgcg gccacaaac tcatcaatgt 720
catcgtcccc gagagccgag cccacttctt ccagcagctg ggctacgtgc tggccacgct 780
gctgctcttc atcctgctac tggtcactgt cctcctggcc gcccgcaggc gccgcggagg 840
ctacgaatac tcggaccaga agtcgggaaa gtcaaagggg aaggatgta acttggcgga 900
gttcgctgtg gctgcagggg accagatgct ttacaggagt gaggacatcc agctagatta 960
caaaaacaac atcctgaagg agagggcgga gctggcccac agccccctgc ctgccaaagta 1020
```

```

catcgacctg gacaaagggg tccggaagga gaactgcaaa tagggaggcc ctgggctcct 1080
ggctgggcca gcagctgcac ctctcctgtc tgtgctcttc ggggcatctc ctgatgctcc 1140
ggggctcacc ccccttccag cggctgggtc cgctttctct gaatttgcc tgggcgtatg 1200
cagaggccgc ctccacaccc ctccccccagg ggcttggtgg cagcatagcc cccaccctg 1260
cggcctttgc tcacgggtgg ccctgcccac ccctggcaca accaaaatcc cactgatgcc 1320
catcatgccc tcagaccctt ctgggctctg cccgctgggg gcctgaagac attcctggag 1380
gacactccca tcagaacctg gcagcccca aactgggggtc agcctcaggg caggagtccc 1440
actcctccag ggctctgtct gtccgggggtc gggagatgtt cctggaggag gacactccca 1500
tcagaacttg gcagccttga agttgggggtc agcctcggca ggagtccac tcctcctggg 1560
gtgctgcctg ccaccaagag ctccccccacc tgtaccacca tgtgggactc caggcaccat 1620
ctgttctccc cagggacctg ctgacttgaa tgccagccct tgctcctctg tgttgctttg 1680
ggccacctgg ggctgcaccc cctgcccctt ctctgcccac tccctaccct agccttgctc 1740
tcagccacct tgatagtcac tgggctccct gtgacttctg accctgacac ccctcccttg 1800
gactctgcct gggctggagt ctagggctgg ggctacattt ggcttctgta ctggctgagg 1860
acaggggagg gagtgaagtt ggtttgggtt ggcctgtgtt gccactctca gcacccaca 1920
tttgcactct ctggtggacc tgccaccatc acaataaagt ccccatctga tttttaaaaa 1980
aaaaaaa

```

<210> 21
 <211> 551
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc feature
 <223> Incyte ID No.: 627722CD1

<400> 21

Met	Glu	Glu	Ala	Glu	Leu	Val	Lys	Gly	Arg	Leu	Gln	Ala	Ile	Thr	1	5	10	15
Asp	Lys	Arg	Lys	Ile	Gln	Glu	Glu	Ile	Ser	Gln	Lys	Arg	Leu	Lys	20	25	30	35
Ile	Glu	Glu	Asp	Lys	Leu	Lys	His	Gln	His	Leu	Lys	Lys	Lys	Ala	40	45	50	55
Leu	Arg	Glu	Lys	Trp	Leu	Leu	Asp	Gly	Ile	Ser	Ser	Gly	Lys	Glu	60	65	70	75
Gln	Glu	Glu	Met	Lys	Lys	Gln	Asn	Gln	Gln	Asp	Gln	His	Gln	Ile	80	85	90	95
Gln	Val	Leu	Glu	Gln	Ser	Ile	Leu	Arg	Leu	Glu	Lys	Glu	Ile	Gln	100	105	110	115
Asp	Leu	Glu	Lys	Ala	Glu	Leu	Gln	Ile	Ser	Thr	Lys	Glu	Glu	Ala	120	125	130	135
Ile	Leu	Lys	Lys	Leu	Lys	Ser	Ile	Glu	Arg	Thr	Thr	Glu	Asp	Ile	140	145	150	155
Ile	Arg	Ser	Val	Lys	Val	Glu	Arg	Glu	Glu	Arg	Ala	Glu	Glu	Ser	160	165	170	175
Ile	Glu	Asp	Ile	Tyr	Ala	Asn	Ile	Pro	Asp	Leu	Pro	Lys	Ser	Tyr	180	185	190	195
Ile	Pro	Ser	Arg	Leu	Arg	Lys	Glu	Ile	Asn	Glu	Glu	Lys	Glu	Asp	200	205	210	215
Asp	Glu	Gln	Asn	Arg	Lys	Ala	Leu	Tyr	Ala	Met	Glu	Ile	Lys	Val	220	225	230	235
Glu	Lys	Asp	Leu	Lys	Thr	Gly	Glu	Ser	Thr	Val	Leu	Ser	Ser	Ile	240	245	250	255
Pro	Leu	Pro	Ser	Asp	Asp	Phe	Lys	Gly	Thr	Gly	Ile	Lys	Val	Tyr				
Asp	Asp	Gly	Gln	Lys	Ser	Val	Tyr	Ala	Val	Ser	Ser	Asn	His	Ser				
Ala	Ala	Tyr	Asn	Gly	Thr	Asp	Gly	Leu	Ala	Pro	Val	Glu	Val	Glu				
Glu	Leu	Leu	Arg	Gln	Ala	Ser	Glu	Arg	Asn	Ser	Lys	Ser	Pro	Thr				

Glu Tyr His Glu	Pro Val Tyr Ala Asn	Pro Phe Tyr Arg Pro Thr	260	265	270
Thr Pro Gln Arg	Glu Thr Val Thr Pro	Gly Pro Asn Phe Gln Glu	275	280	285
Arg Ile Lys Ile	Lys Thr Asn Gly Leu	Gly Ile Gly Val Asn Glu	290	295	300
Ser Ile His Asn	Met Gly Asn Gly Leu	Ser Glu Glu Arg Gly Asn	305	310	315
Asn Phe Asn His	Ile Ser Pro Ile Pro	Pro Val Pro His Pro Arg	320	325	330
Ser Val Ile Gln	Gln Ala Glu Glu Lys	Leu His Thr Pro Gln Lys	335	340	345
Arg Leu Met Thr	Pro Trp Glu Glu Ser	Asn Val Met Gln Asp Lys	350	355	360
Asp Ala Pro Ser	Pro Lys Pro Arg Leu	Ser Pro Arg Glu Thr Ile	365	370	375
Phe Gly Lys Ser	Glu His Gln Asn Ser	Ser Pro Thr Cys Gln Glu	380	385	390
Asp Glu Glu Asp	Val Arg Tyr Asn Ile	Val His Ser Leu Pro Pro	395	400	405
Asp Ile Asn Asp	Thr Glu Pro Val Thr	Met Ile Phe Met Gly Tyr	410	415	420
Gln Gln Ala Glu	Asp Ser Glu Glu Asp	Lys Lys Phe Leu Thr Gly	425	430	435
Tyr Asp Gly Ile	Ile His Ala Glu Leu	Val Val Ile Asp Asp Glu	440	445	450
Glu Glu Glu Asp	Glu Gly Glu Ala Glu	Lys Pro Ser Tyr His Pro	455	460	465
Ile Ala Pro His	Ser Gln Val Tyr Gln	Pro Ala Lys Pro Thr Pro	470	475	480
Leu Pro Arg Lys	Arg Ser Glu Ala Ser	Pro His Glu Asn Thr Asn	485	490	495
His Lys Ser Pro	His Lys Asn Ser Ile	Ser Leu Lys Glu Gln Glu	500	505	510
Glu Ser Leu Gly	Ser Pro Val His His	Ser Pro Phe Asp Ala Gln	515	520	525
Thr Thr Gly Asp	Gly Thr Glu Asp Pro	Ser Leu Thr Ala Leu Arg	530	535	540
Met Arg Met Ala	Lys Leu Gly Lys Lys	Val Ile	545	550	

<210> 22

<211> 99

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No.: 1556751CD1

<400> 22

Met Glu Ala Leu Ala	Asn Val Asn Phe	Pro Arg Lys Ser Phe Arg	1	5	10	15
Pro Glu Asp Ala Gly	Lys Glu Ser Gly	Ser Gln Gly Gly Phe Cys	20	25	30	
Val Pro Ala Ala Arg	Pro Gln Thr Met	Val Thr Gly Pro Ser Cys	35	40	45	
Ser Ser Pro Gly Leu	Gln Asn Phe Ser	Pro Gln Arg Lys Glu Asn	50	55	60	
Arg Ala Cys Ala Cys	Trp Gln Asn Ala	Gly Pro Ala Pro Lys Asn	65	70	75	
Pro Met Cys Val Arg	Leu Lys Val Gly	Arg Pro Gln Ala Ser Gln				

80
Arg Lys Leu Lys Glu Thr Gly Leu Cys
95

85

90

<210> 23
<211> 493
<212> PRT
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No.: 2268890CD1

<400> 23

Met	Arg	Pro	Leu	Cys	Val	Thr	Cys	Trp	Trp	Leu	Gly	Leu	Leu	Ala	15
1				5					10						
Ala	Met	Gly	Ala	Val	Ala	Gly	Gln	Glu	Asp	Gly	Phe	Glu	Gly	Thr	30
				20					25						
Glu	Glu	Gly	Ser	Pro	Arg	Glu	Phe	Ile	Tyr	Leu	Asn	Arg	Tyr	Lys	45
				35					40						
Arg	Ala	Gly	Glu	Ser	Gln	Asp	Lys	Cys	Thr	Tyr	Thr	Phe	Ile	Val	60
				50					55						
Pro	Gln	Gln	Arg	Val	Thr	Gly	Ala	Ile	Cys	Val	Asn	Ser	Lys	Glu	75
				65					70						
Pro	Glu	Val	Leu	Leu	Glu	Asn	Arg	Val	His	Lys	Gln	Glu	Leu	Glu	90
				80					85						
Leu	Leu	Asn	Asn	Glu	Leu	Leu	Lys	Gln	Lys	Arg	Gln	Ile	Glu	Thr	105
				95					100						
Leu	Gln	Gln	Leu	Val	Glu	Val	Asp	Gly	Gly	Ile	Val	Ser	Glu	Val	120
				110					115						
Lys	Leu	Leu	Arg	Lys	Glu	Ser	Arg	Asn	Met	Asn	Ser	Arg	Val	Thr	135
				125					130						
Gln	Leu	Tyr	Met	Gln	Leu	Leu	His	Glu	Ile	Ile	Arg	Lys	Arg	Asp	150
				140					145						
Asn	Ala	Leu	Glu	Leu	Ser	Gln	Leu	Glu	Asn	Arg	Ile	Leu	Asn	Gln	165
				155					160						
Thr	Ala	Asp	Met	Leu	Gln	Leu	Ala	Ser	Lys	Tyr	Lys	Asp	Leu	Glu	180
				170					175						
His	Lys	Tyr	Gln	His	Leu	Ala	Thr	Leu	Ala	His	Asn	Gln	Ser	Glu	195
				185					190						
Ile	Ile	Ala	Gln	Leu	Glu	Glu	His	Cys	Gln	Arg	Val	Pro	Ser	Ala	210
				200					205						
Arg	Pro	Val	Pro	Gln	Pro	Pro	Pro	Ala	Ala	Pro	Pro	Arg	Val	Tyr	225
				215					220						
Gln	Pro	Pro	Thr	Tyr	Asn	Arg	Ile	Ile	Asn	Gln	Ile	Ser	Thr	Asn	240
				230					235						
Glu	Ile	Gln	Ser	Asp	Gln	Asn	Leu	Lys	Val	Leu	Pro	Pro	Pro	Leu	255
				245					250						
Pro	Thr	Met	Pro	Thr	Leu	Thr	Ser	Leu	Pro	Ser	Ser	Thr	Asp	Lys	270
				260					265						
Pro	Ser	Gly	Pro	Trp	Arg	Asp	Cys	Leu	Gln	Ala	Leu	Glu	Asp	Gly	285
				275					280						
His	Asp	Thr	Ser	Ser	Ile	Tyr	Leu	Val	Lys	Pro	Glu	Asn	Thr	Asn	300
				290					295						
Arg	Leu	Met	Gln	Val	Trp	Cys	Asp	Gln	Arg	His	Asp	Pro	Gly	Gly	315
				305					310						
Trp	Thr	Val	Ile	Gln	Arg	Arg	Leu	Asp	Gly	Ser	Val	Asn	Phe	Phe	330
				320					325						
Arg	Asn	Trp	Glu	Thr	Tyr	Lys	Gln	Gly	Phe	Gly	Asn	Ile	Asp	Gly	345
				335					340						
Glu	Tyr	Trp	Leu	Gly	Leu	Glu	Asn	Ile	Tyr	Trp	Leu	Thr	Asn	Gln	360
				350					355						

Gly	Asn	Tyr	Lys	Leu	Leu	Val	Thr	Met	Glu	Asp	Trp	Ser	Gly	Arg	
				365					370					375	
Lys	Val	Phe	Ala	Glu	Tyr	Ala	Ser	Phe	Arg	Leu	Glu	Pro	Glu	Ser	
				380					385					390	
Glu	Tyr	Tyr	Lys	Leu	Arg	Leu	Gly	Arg	Tyr	His	Gly	Asn	Ala	Gly	
				395					400					405	
Asp	Ser	Phe	Thr	Trp	His	Asn	Gly	Lys	Gln	Phe	Thr	Thr	Leu	Asp	
				410					415					420	
Arg	Asp	His	Asp	Val	Tyr	Thr	Gly	Asn	Cys	Ala	His	Tyr	Gln	Lys	
				425					430					435	
Gly	Gly	Trp	Trp	Tyr	Asn	Ala	Cys	Ala	His	Ser	Asn	Leu	Asn	Gly	
				440					445					450	
Val	Trp	Tyr	Arg	Gly	Gly	His	Tyr	Arg	Ser	Arg	Tyr	Gln	Asp	Gly	
				455					460					465	
Val	Tyr	Trp	Ala	Glu	Phe	Arg	Gly	Gly	Ser	Tyr	Ser	Leu	Lys	Lys	
				470					475					480	
Val	Val	Met	Met	Ile	Arg	Pro	Asn	Pro	Asn	Thr	Phe	His			
				485					490						